

SYMMETRIC PAIRS AND GORELIK ELEMENTS

MICHEL DUFLO AND EMANUELA PETRACCI

To Ernest Borisovich Vinberg on the occasion of his seventieth birthday

ABSTRACT. The second author gave a formula for the elements of the enveloping algebra of a Lie superalgebra defined by Gorelik under an appropriate unimodularity assumption. We show that this formula is a particular case of a formula for the Jacobian of the exponential map of a symmetric superspace.

CONTENTS

Introduction.	1
1. Berezinians and dualizing modules.	4
1.1. Notations for modules.	4
1.2. Symmetric algebras as coalgebras.	5
1.3. Functional calculus.	6
1.4. Berezinians and supertraces.	7
1.5. Dualizing modules.	9
1.6. Dualizing modules and duality.	11
2. Dualizing modules on formal Lie supergroups.	11
2.1. Enveloping algebras as coalgebras.	12
2.2. The exponential map of a formal Lie supergroup.	13
2.3. The generic point of a Lie superalgebra.	13
2.4. The Jacobian of the exponential map.	15
3. Formal homogeneous spaces.	16
3.1. Induced and coinduced representations in exponential coordinates.	16
3.2. The exponential map on a formal homogeneous space	17
3.3. Imprimitivity theorem.	18
3.4. Dualizing modules on formal homogeneous spaces.	18
3.5. The Jacobian of the exponential map on a formal homogeneous space.	19
3.6. Homogeneous spaces of purely odd dimension.	20
4. Symmetric pairs.	22
4.1. Representations by coderivations.	22
4.2. Induced and coinduced representations.	25
4.3. The Jacobian of the exponential map for a symmetric pair.	28
4.4. Gorelik elements.	30
References	31

INTRODUCTION.

In her PhD thesis [17], the second author gave a formula for the elements, in the enveloping algebra of a Lie superalgebra, defined by Gorelik [9] under an appropriate unimodularity assumption. We recall Gorelik's definition. Let $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ be a finite dimensional superalgebra over a field of characteristic 0, and σ be the automorphism of $\mathfrak{g}$ such that $\sigma(a) = a$ if $a \in \mathfrak{g}_0$

1991 *Mathematics Subject Classification.* 17B35, 14L05, 22E60.

Key words and phrases. Lie algebra, Lie superalgebra, exponential map, Jacobian, Casimir ghost, Gorelik element, Lie supergroup, formal Lie supergroup.

The research of the second author was done while she was working at the Mathematics department *Section de mathématiques* of the University of Geneva and received some financial support from the Swiss National Science Foundation. During part of the final redaction she was supported first by a post-doctoral grant of the European network LIEGRITS at the I.E.C.N. Institute in Nancy, and then by an invitation at the I.H.E.S Institute in Bures-sur-Yvettes. The second author thanks all these institutions. A special thanks to Prof. Caroline Gruson, LIEGRITS responsible in Nancy.

and $\sigma(a) = -a$ if $a \in \mathfrak{g}_1$. Arnaudon, Bauer and Frappat [1] define the *twisted adjoint action* $\text{ad}'(a)$ of $a \in \mathfrak{g}$ in the enveloping algebra $U(\mathfrak{g})$ by

$$(1) \quad \text{ad}'(a)(u) = au - \pm u\sigma(a) \quad \text{for} \quad u \in U(\mathfrak{g}).$$

Let $S(\mathfrak{g})$ be the symmetric algebra and $\beta : S(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ be the symmetrization map. The finite dimensional¹ subspace $\beta(S(\mathfrak{g}_1))$ of $U(\mathfrak{g})$ is stable under the twisted adjoint action (see [1]). Gorelik [9] proves that *if the following unimodularity condition holds,*

$$(2) \quad \text{str}_{\mathfrak{g}_1}(\text{ad } a) = 0 \quad \text{for} \quad a \in \mathfrak{g}_0,$$

there exists a non zero invariant of the twisted adjoint action $T \in \beta(S(\mathfrak{g}_1))$, unique up to a multiplicative constant. We call T a *Gorelik element* of $U(\mathfrak{g})$. The Gorelik elements occurred already for $\mathfrak{osp}(1, 2)$ in [14, 19], for $\mathfrak{osp}(1, 2l)$ in [1] under the name *Casimir ghosts*, and in [16]. The Gorelik elements play an important role in the representation theory of $\mathfrak{g}$, see in particular [9–11, 16, 19].

Formula for Gorelik's elements. Let us describe the formula given in [17] for the Gorelik elements. Let $q = \dim \mathfrak{g}_1$. Choose a basis d of the 1-dimensional space $S^q(\mathfrak{g}_1)$. Consider the algebra $S(\mathfrak{g}_1)^*$ dual of the coalgebra $S(\mathfrak{g}_1)$. By duality, $S(\mathfrak{g}_1)$ is a module over $S(\mathfrak{g}_1)^*$; this action is the usual interior product and the map $f \rightarrow fd$ is an isomorphism from $S(\mathfrak{g}_1)^*$ to $S(\mathfrak{g}_1)$. In particular, to describe our formula means to describe a special element $J_2 \in S(\mathfrak{g}_1)^*$. To do this, we use the *generic point of $\mathfrak{g}_1$* . It is the element $y = \sum e_i x^i$ of the Lie $S(\mathfrak{g}_1)^*$ -superalgebra $\mathfrak{g} \otimes S(\mathfrak{g}_1)^*$, where $(e_i)_{1 \leq i \leq q}$ is a basis of $\mathfrak{g}_1$ and $(x^i)_{1 \leq i \leq q}$ is the corresponding dual basis of $\mathfrak{g}_1^*$. For $k \in \mathbb{N}$, we have

$$(3) \quad \text{ad}^k y = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq q} \sum_{s \in S_k} \pm \text{ad } e_{i_{s(1)}} \cdots \text{ad } e_{i_{s(k)}} x^{i_1} \cdots x^{i_k}.$$

In particular, $\text{ad}^k y = 0$ if $k > q$. When k is even the space $\mathfrak{g}_1$ is invariant by $\text{ad } e_{i_{s(1)}} \cdots \text{ad } e_{i_{s(k)}}$. We consider the finite sum

$$(4) \quad \frac{\text{sh}(\text{ad } \frac{y}{2})}{\text{ad } \frac{y}{2}} = 1 + \frac{\text{ad}^2 y}{24} + \frac{\text{ad}^4 y}{1920} + \dots$$

It is an even invertible endomorphism of $\mathfrak{g} \otimes S(\mathfrak{g}_1)^*$ which stabilizes $\mathfrak{g}_1 \otimes S(\mathfrak{g}_1)^*$. We denote by $J_2 \in S(\mathfrak{g}_1)^*$ the Berezinian (in this case the inverse of the determinant) of this endomorphism of $\mathfrak{g}_1 \otimes S(\mathfrak{g}_1)^*$. We summarize this definition by the notation

$$(5) \quad J_2 = \text{Ber}_{\mathfrak{g}_1} \left(\frac{\text{sh}(\text{ad } \frac{y}{2})}{\text{ad } \frac{y}{2}} \right).$$

Similarly, if k is even, we define

$$(6) \quad \text{str}_{\mathfrak{g}_1}(\text{ad}^k y) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq q} \sum_{s \in S_k} \pm \text{str}_{\mathfrak{g}_1}(\text{ad } e_{i_{s(1)}} \cdots \text{ad } e_{i_{s(k)}}) x^{i_1} \cdots x^{i_k} \in S(\mathfrak{g}_1)^*.$$

The relation $\text{Ber}(\exp(\cdot)) = \exp(\text{str}(\cdot))$ allows to write explicitly the Taylor expansion of J_2 in homogeneous components:

$$(7) \quad J_2 = 1 + \frac{1}{24} \text{str}_{\mathfrak{g}_1}(\text{ad}^2 y) - \frac{\text{str}_{\mathfrak{g}_1}(\text{ad}^4 y)}{2880} + \frac{\text{str}_{\mathfrak{g}_1}^2(\text{ad}^2 y)}{1152} + \dots$$

For instance, for $q = 2$ we obtain

$$(8) \quad J_2 = 1 + \frac{1}{24} \text{str}_{\mathfrak{g}_1}(-\text{ad } e_1 \text{ad } e_2 + \text{ad } e_2 \text{ad } e_1) x^1 x^2.$$

We have —see [17] and Corollary 4.3 below:

Theorem 0.1. *Under the unimodularity condition (2), $\beta(J_2 d)$ is a Gorelik element of $U(\mathfrak{g})$.*

¹Since $\mathfrak{g}_1$ is purely odd, $S(\mathfrak{g}_1)$ is isomorphic to the exterior algebra of $\mathfrak{g}_1$ (see remark 1.1).

For instance, for $q = 2$, if we choose $d = e_1 e_2$ we obtain the Gorelik element

$$(9) \quad T = \frac{1}{2}(j(e_1)j(e_2) - j(e_2)j(e_1)) + \frac{1}{24} \text{str}_{\mathfrak{g}_1}(\text{ad } e_1 \text{ ad } e_2 - \text{ad } e_2 \text{ ad } e_1),$$

where $j : \mathfrak{g} \rightarrow U(\mathfrak{g})$ is the canonical injection.

Occurrence of formal symmetric spaces. The occurrence of the operator $\frac{\text{sh}(\text{ad } \frac{y}{2})}{\text{ad } \frac{y}{2}}$ suggests a relation with the differential of the exponential map. Indeed, *we prove in this paper that Theorem 0.1 is a particular case of a formula for the Jacobian of the exponential map for symmetric spaces —in the sense of supergeometry.* A natural setting to present this result is to consider a Lie superalgebra $\mathfrak{g}$ over a commutative $\mathbb{Q}$ -superalgebra $\mathbb{K}$, and a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ such that the quotient $\mathfrak{g}/\mathfrak{h}$ is free of finite rank over $\mathbb{K}$. The formal Lie supergroups G , H and the formal homogeneous space G/H can be defined by mean of suitable commutative $\mathbb{K}$ -superalgebras $F(G)$, $F(H)$, $F(G/H)$ of formal functions. Formal supergroups were first considered by Berezin and Kac [3]. There exists and we choose a formal diffeomorphism $\theta : \mathfrak{g}/\mathfrak{h} \rightarrow G/H$, or equivalently an algebra isomorphism $\theta^* : F(G/H) \rightarrow F(\mathfrak{g}/\mathfrak{h}) := S(\mathfrak{g}/\mathfrak{h})^*$ — i.e. into the algebra of formal “functions” on $\mathfrak{g}/\mathfrak{h}$, which is an algebra of formal power series in a finite number of even and odd variables. In this situation, the *dualizing* $F(G/H)$ -module $K(F(G/H))$ is defined. It is free of rank 1: in the purely even case (i.e. $\mathbb{K} = \mathbb{K}_0$ and $\mathfrak{g} = \mathfrak{g}_0$) it is the module of top degree differential forms on G/H , and in general it is the Berezinian of the module of 1-differential forms (see [25] and subsection 1.5 below). There is a natural action of $\mathfrak{g}$ in $K(F(G/H))$ by Lie derivatives, and a natural $\mathbb{K}$ -linear isomorphism $\theta^* : K(F(G/H)) \rightarrow K(F(\mathfrak{g}/\mathfrak{h}))$. The unimodularity assumption reads:

$$(10) \quad \text{str}_{\mathfrak{g}/\mathfrak{h}}(\text{ad } a) = 0 \quad \text{for} \quad a \in \mathfrak{h}.$$

We show in proposition 3.3 that condition (10) is equivalent to the existence of a $\mathfrak{g}$ -invariant basis $D_{G/H} \in K(F(G/H))$. In this case, there is a translation invariant element $d_{\mathfrak{g}/\mathfrak{h}} \in K(F(\mathfrak{g}/\mathfrak{h}))$ and a function $J \in F(\mathfrak{g}/\mathfrak{h})$, the *Jacobian* of θ , such that

$$(11) \quad \theta^*(D_{G/H}) = J d_{\mathfrak{g}/\mathfrak{h}} \text{ and } J(0) = 1.$$

We compute this Jacobian in two cases.

First, we suppose that $\mathfrak{h} = 0$ and that $\mathfrak{g}$ is a free $\mathbb{K}$ -module of finite rank. In theorem 2.1 we give a formula for the Jacobian of the exponential map $\mathfrak{g} \rightarrow G$. The result is not surprising, but we think that it is a good introduction to the second case.

In the second case, we assume that $\mathfrak{h}$ is the fixed point set of an involution σ of $\mathfrak{g}$. We write $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$ the decomposition of $\mathfrak{g}$ in 1 and -1 eigenspaces of σ . We say that $(\mathfrak{g}, \sigma)$ or $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$ is a symmetric pair, and that G/H is symmetric. The exponential map is a formal diffeomorphism $\mathfrak{q} \rightarrow G/H$. We prove (see theorem 4.6 below):

Theorem 0.2. *Suppose that $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$ is a symmetric pair, that $\mathfrak{q}$ is free of finite rank, and that condition (10) holds. Then the Jacobian of the exponential map $\mathfrak{q} \rightarrow G/H$ is the element $J \in F(\mathfrak{q})$ given by the formula*

$$(12) \quad J = \text{Ber}_{\mathfrak{q}} \left(\frac{\text{sh}(\text{ad } y)}{\text{ad } y} \right),$$

where y is the generic point of $\mathfrak{q}$.

We illustrate Theorem 0.2 in two particular extreme cases.

Example 0.1. *Suppose that $\mathbb{K} = \mathbb{R}$ and that $\mathfrak{g} = \mathfrak{g}_0$ is a finite dimensional Lie algebra. Let $\mathbf{G}$ and $\mathbf{H}$ be connected Lie groups with Lie algebras $\mathfrak{g}$ and $\mathfrak{h}$, and such that $\mathbf{H}$ is a closed subgroup of $\mathbf{G}$. Condition (10) is equivalent to the existence of a non zero left invariant differential form of maximal degree $D_{\mathbf{G}/\mathbf{H}}$ on $\mathbf{G}/\mathbf{H}$. It is well known (see for instance [21]) that its inverse image*

by the exponential map $\exp : \mathfrak{q} \rightarrow \mathbf{G}/\mathbf{H}$ at a point $a \in \mathfrak{q}$ is equal to $\det_{\mathfrak{q}}(\frac{\text{sh}(\text{ad } a)}{\text{ad } a})d_{\mathfrak{q}}$, where $d_{\mathfrak{q}}$ is a form of maximal degree on $\mathfrak{q}$ invariant by translations. Since in this case $\text{Ber}_{\mathfrak{q}} = \det_{\mathfrak{q}}$, we recover Theorem 0.2 by taking the Taylor series of $\det_{\mathfrak{q}}(\frac{\text{sh}(\text{ad } a)}{\text{ad } a})$ at $a = 0$.

Example 0.2. Suppose that $\mathbb{K} = \mathbb{K}_0$ is a field of characteristic 0, and consider a finite dimensional Lie superalgebra $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$. This is a symmetric pair, and condition (10) is the same than Gorelik's condition (2). Assuming this condition, we prove in theorem 3.1 a result on induced representations of a Lie superalgebra which says in particular that the image $\tilde{\beta}(Jd)$ of $\beta(Jd)$ in $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{g}_0$ is a non zero $\mathfrak{g}$ -invariant element. We explain in subsection 4.4 how Theorem 0.1 follows by considering an isomorphism of $\mathfrak{g}$ -modules from $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{g}_0$ to the subspace $\beta(S(\mathfrak{g}_1))$ of $U(\mathfrak{g})$ endowed with the twisted adjoint action.

In fact, it is better not to assume condition (10): in general, we twist $K(F(G/H))$ by a suitable line bundle on G/H to get the existence of an invariant element. More precisely, we consider the $\mathfrak{g}$ -module $F(G/H, \text{Ber } \mathfrak{g}/\mathfrak{h})$ coinduced by the rank 1 $\mathfrak{h}$ -module provided by the character $\text{str}_{\mathfrak{g}/\mathfrak{h}}$ (for the details of these notations, see section 3). It is a free $F(G/H)$ -module of rank 1, as well as the *twisted dualizing module* $K(F(G/H), \text{Ber } \mathfrak{g}/\mathfrak{h}) := K(F(G/H)) \otimes_{F(G/H)} F(G/H, \text{Ber } \mathfrak{g}/\mathfrak{h})$. We prove in proposition 3.4 that there is a canonical $\mathfrak{g}$ -invariant basis $\mathbf{D}_{G/H} \in K(F(G/H), \text{Ber } \mathfrak{g}/\mathfrak{h})$. In Theorem 4.6, we prove formula (12) also in the twisted case.

Formulas related to the symmetrization map. Our proofs involve many useful formulas about the symmetrization β , in particular how to work with induced and coinduced representations in exponential coordinates. We think that they are of interest, independently of the present application to Jacobians. In the Lie group case $\mathbf{G}/\mathbf{H}$ these formulas occur more or less explicitly in Rouvière [21, 22]. In the case of the symmetric pair $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ associated to a finite dimensional Lie superalgebra defined over a field of characteristic zero, they are given in Koszul [13]. However, our proofs are different. We use the methods of [17, 18]: we seek *universal formulas*, that is formulas which can be expressed in the same manner for all symmetric pairs $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$. They depend on a finite number of formal power series in $\mathbb{K}_0[[t]]$ which must satisfy some *functional equations*. We are lucky enough: these functional equations have solutions which are essentially unique, and which are given by explicit formulas.

Hypotheses used in the text. We assume from the beginning that $\mathbb{K}$ contains $\mathbb{Q}$, because the main actors of this paper are the symmetrization $\beta : S(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ and the related exponential map $\mathfrak{g} \rightarrow G$. But nothing more. Besides the obvious fact that it is better to avoid unnecessary hypotheses, adding more assumptions on $\mathbb{K}$ (like $\mathbb{K}$ being a field, or the weaker assumption $\mathbb{K} = \mathbb{K}_0$) would not be specially helpful, because, as we have seen in formula (5), it is convenient to use Lie superalgebras over rings like exterior algebras.

For simplicity, we state and prove our results involving Berezinians and friends when $\mathfrak{g}/\mathfrak{h}$ is free of finite rank, even if they can probably easily be extended to the case when $\mathfrak{g}/\mathfrak{h}$ is projective of finite rank, or to a locally free sheaf version of it. For the other results, we suppose that $\mathfrak{h}$ is a direct factor of $\mathfrak{g}$, but we do not require anything more on the $\mathbb{K}$ -module structure of $\mathfrak{g}$ or $\mathfrak{h}$.

Working in this generality forces us to write more natural proofs. The drawback is that we have to use some results usually stated under the assumption that $\mathbb{K}$ is a field. The main one is the Poincaré-Birkhoff-Witt theorem which states that the symmetrization $\beta : S(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ is an isomorphism of coalgebras. This one is proved in [8] –see also [18]. For some others, like Blattner's imprimitivity theorem [4], we provide the proofs.

1. BEREZINIANS AND DUALIZING MODULES.

1.1. Notations for modules. In this text, *all $\mathbb{Z}$ -modules are supposed to be $\mathbb{Z}/2\mathbb{Z}$ -graded*. Let $M = M_0 \oplus M_1$ be a $\mathbb{Z}$ -module. The elements of M_0 are called even, and the elements of M_1

odd. We denote by $p(m) \in \mathbb{Z}/2\mathbb{Z} = \{0, 1\}$ the parity of an element m , in the following sense: whenever the expression $p(m)$ occurs in a formula, it is implicitly assumed that $m \in M_{p(m)}$.

Depending on the context, we use the notations id_M , id or 1 for the identity map of M . We write Π for the change of parity functor. Thus $(\Pi M)_0 = M_1$, $(\Pi M)_1 = M_0$. We use the notations π_M or π for the identity map of M when it is considered as a map from $M \rightarrow \Pi M$.

We fix a commutative $\mathbb{Q}$ -superalgebra $\mathbb{K} = \mathbb{K}_0 \oplus \mathbb{K}_1$, associative with unit $1 \in \mathbb{K}_0$ (supercommutative in the terminology of [15, ch. 3]).

If $M = M_0 \oplus M_1$ is a $\mathbb{K}$ -module, we write $km = \pm mk$ for $k \in \mathbb{K}$ and $m \in M$, where here and after, the symbols are implicitly assumed to be homogeneous, and the sign $\pm$ is obtained by the standard rule of signs (see [15, ch. 3.1]). For instance in this case, $\pm = (-1)^{p(k)p(m)}$.

If M and N are $\mathbb{K}$ -modules, if $\mathbb{K}$ is clear from the context we write $M \otimes N$ for $M \otimes_{\mathbb{K}} N$, $\text{Hom}(M, N)$ for $\text{Hom}_{\mathbb{K}}(M, N)$, $\text{End}(M)$ for $\text{Hom}(M, M)$, M^* for $\text{Hom}(M, \mathbb{K})$. We recall that the $\mathbb{K}$ -structure on ΠM is defined by $\pi_M(m)k = \pi_M(mk)$, i.e. by the requirement $\pi_M \in \text{Hom}(M, \Pi M)_1$.

Let M be a free $\mathbb{K}$ -module of finite rank. It is isomorphic to $\mathbb{K}^p \oplus \Pi \mathbb{K}^q$ for some uniquely determined p and q in $\mathbb{N}$. We say that $p + q \in \mathbb{N}$ is the rank of M . If we need to keep track of parities, we say that the rank is $(p, q) \in \mathbb{N}^2$. For instance, a module of rank 1 is isomorphic either to $\mathbb{K}$ (in this case it is of rank $(1, 0)$) or to $\Pi \mathbb{K}$ (in this case it is of rank $(0, 1)$).

Suppose $(e_i)_{i \in I}$ is a basis of a free module M of finite rank. In this text we always suppose that basis are homogeneous. We denote by $(x^i)_{i \in I}$ the dual basis of M^* : $x^i(e_j) = \delta_j^i$. If N is a $\mathbb{K}$ -module, we identify $\text{Hom}(M, N)$ and $N \otimes M^*$ in the usual manner. This identification sends $\text{id}_M \in \text{End}(M)_0$ to $\sum e_i \otimes x^i$, and $\pi_M \in \text{Hom}(M, \Pi M)_1$ to $\sum \pi_M e_i \otimes x^i$.

1.2. Symmetric algebras as coalgebras. In this subsection, we state the properties of formal functions on a $\mathbb{K}$ -module which we shall need in the sequel. For a more detailed exposition, we refer to [18, 20]. Let M be a $\mathbb{K}$ -module. We denote by $S(M) = \bigoplus_{n \geq 0} S^n(M)$ the symmetric algebra of M . It is an Hopf superalgebra.

Remark 1.1. *Everything in this article is considered in the super setting. Thus, $S(M)$ is the commutative superalgebra freely generated by M . If M is a free $\mathbb{K}$ -module with basis $(e_i)_{1 \leq i \leq n}$, $S(M)$ is equal to the commutative superalgebra $\mathbb{K}[e_1, \dots, e_n]$ of finite sums with coefficients in $\mathbb{K}$ of monomials $e_1^{p_1} \dots e_n^{p_n}$, with $p_i \leq 1$ if e_i is odd.*

In particular, if $(e_1, \dots, e_q)$ is a basis of M consisting of q odd elements, $S(M) = \mathbb{K}[e_1, \dots, e_q]$ is the exterior algebra on the generators e_i . It is a free $\mathbb{K}$ -module of rank 2^q .

We denote by $\Delta_{S(M)} \in \text{Hom}(S(M), S(M) \otimes S(M))$ the coproduct, and by $S_{S(M)} : S(M) \rightarrow S(M)$ the antipode. Let $w \in S(M)$, we denote its coproduct by

$$\Delta_{S(M)}(w) = \sum w_i \otimes w'_i.$$

We denote by $F(M)$ the algebra $S(M)^*$ dual to the coalgebra $S(M)$, and we consider it as the algebra of formal $\mathbb{K}$ -valued functions defined in a formal neighborhood of $0 \in M$. We denote by $\delta : F(M) \rightarrow \mathbb{K}$ the map defined by $\delta(f) = f(1)$. It is a morphism of $\mathbb{K}$ -superalgebras, and we consider it as the “evaluation at $0 \in M$ ”. Let $S^+(M) = \ker \delta$. Thus $S^+(M) = \bigoplus_{n > 0} S^n(M)$.

Let N be a $\mathbb{K}$ -module. We denote by $F(M, N) = \text{Hom}(S(M), N)$ the space of formal N -valued functions on M . We identify N with the submodule of $F(M, N)$ consisting of the “constant functions”, that is the functions which are zero over $S^+(M)$. We still denote by $\delta : F(M, N) \rightarrow N$ the map defined by $\delta(f) = f(1)$.

The space $F(M, N)$ is an $F(M)$ -module. More generally, if $\mu : N_1 \otimes N_2 \rightarrow N$ is a map of $\mathbb{K}$ -modules, there is a corresponding map $\tilde{\mu} : F(M, N_1) \otimes_{F(M)} F(M, N_2) \rightarrow F(M, N)$, defined by the formula

$$(13) \quad \tilde{\mu}(f_1 \otimes f_2) = \mu \circ (f_1 \otimes f_2) \circ \Delta_{S(M)}, \quad \text{for } f_1 \in F(M, N_1), f_2 \in F(M, N_2).$$

Depending on the setting, there are various notations for $\tilde{\mu}(f_1 \otimes f_2)$. For instance, if N is a Lie $\mathbb{K}$ -superalgebra with product $[n_1, n_2] = \mu(n_1 \otimes n_2)$, we usually employ the notation $[f_1, f_2] = \tilde{\mu}(f_1 \otimes f_2)$, which gives to $F(M, N)$ a Lie $F(M)$ -superalgebra structure. Similarly, if N is an associative algebra with product $n_1 n_2 = \mu(n_1 \otimes n_2)$, we usually employ the notation $f_1 f_2 = \tilde{\mu}(f_1 \otimes f_2)$.

However, in the special case $N = S(M)$ we also use the notation $f_1 * f_2 = \tilde{\mu}(f_1 \otimes f_2)$. Indeed, it is the traditional notation (see e.g. [20]), and it helps to avoid confusions with the composition of maps in $F(M, S(M)) = \text{End}(S(M))$, which is a different associative product. The constant function $1 \in F(M, S(M))$ is the unit for the product $*$. The identity $\text{id}_{S(M)} \in \text{End}_{\mathbb{K}}(S(M))$ and the antipode $S_{S(M)} \in \text{End}_{\mathbb{K}}(S(M))$ are mutual inverses for the product $*$.

Let $\phi \in \text{End}_{\mathbb{K}}(S(M))$. We use the notation $c_\phi = \phi * \text{id}_{S(M)}$ and we denote by ζ_ϕ its transpose acting (as a derivation) in $F(M, N)$. For $w \in S(M)$ and $g \in F(M, N)$, we have:

$$(14) \quad c_\phi(w) = \sum \phi(w_i) w'_i \text{ and } \zeta_\phi(g) = \pm g \circ c_\phi.$$

Note that any $\psi \in \text{End}_{\mathbb{K}}(S(M))$ is of the form $\psi = c_\phi$ with $\phi = S_{S(M)} * \psi$. We consider particular cases.

- i) Let $f \in F(M)$. Then $\zeta_f \in \text{End}_{\mathbb{K}}(F(M, N))$ is the left multiplication by f which defines the $F(M)$ -module structure of $F(M, N)$. Thus we just write $fg = \zeta_f(g)$ for $g \in F(M, N)$. Accordingly, for $w \in S(M)$, we also write fw for $c_f(w)$. *This provides $S(M)$ with a structure of $F(M)$ -module.*

Note that if $f \in M^*$, then c_f is a derivation of $S(M)$, it is the derivative in the direction f . So, for a general $f \in F(M)$, c_f may be thought of as an infinite order differential operator with constant coefficients on $S(M)$. However, in this paper, we prefer to consider $c_f(w)$ as “the product of the distribution w by the function f ”.

- ii) Let $w \in S(M)$. Then c_w is the left multiplication by w . Thus, for $f \in F(M, N)$ and $w' \in S(M)$ we have:

$$(15) \quad \zeta_w(f)(w') = \pm f(w w').$$

We consider ζ_w as a constant coefficient differential operator on $F(M, N)$. In particular, for $m \in M$, ζ_m is the derivative in the direction m .

The space $F(M, M) \subset \text{End}(S(M))$ is called the space of *formal vector fields* on M . Let $\alpha \in F(M, M)$. In this case c_α is a coderivation of $S(M)$ and ζ_α a derivation of $F(M)$. For α and β in $F(M, M)$, we have (see [18, 20]):

$$(16) \quad [c_\alpha, c_\beta] = -c_\gamma \text{ and } [\zeta_\alpha, \zeta_\beta] = \zeta_\gamma, \text{ where } \gamma = \zeta_\alpha(\beta) - \pm \zeta_\beta(\alpha).$$

We denote by x_M the identity map of M , considered as the element of $F(M, M)$ which is 0 on $S^n(M)$ for $n \neq 1$. We call it the *generic point* of M .

1.3. Functional calculus. Let M and N be $\mathbb{K}$ -modules. We say that a family $(f_i)_{i \in I}$ of elements of $F(M, N)$ is locally finite if, for all $w \in S(M)$, the number of elements $i \in I$ such that $f_i(w) \neq 0$ is finite. In this case, we denote by

$$(17) \quad \sum_{i \in I} f_i \in F(M, N)$$

the element such that

$$(18) \quad \left(\sum_{i \in I} f_i \right)(w) = \sum_{i \in I} f_i(w) \text{ for } w \in S(M).$$

Let A be an associative $\mathbb{K}$ -superalgebra with unit $1 \in A_0$, and M be a $\mathbb{K}$ -module. Let us consider an $X \in F(M, A)_0$ such that $\delta(X) = 0$. For $w \in S^k(M)$, we have $X^n(w) = 0$ for $n > k$.

Thus, for $f = \sum_{n \geq 0} f_n t^n \in \mathbb{K}_0[[t]]$ we can define $f(X) \in F(M, A)_0$ by the locally finite sum

$$(19) \quad f(X) = \sum_{n \geq 0} f_n X^n.$$

The rules of functional calculus apply: $f \mapsto f(X)$ is a $\mathbb{K}_0$ -algebra homomorphism, and $\delta(f(X)) = f_0$. Moreover, if $g \in \mathbb{K}_0[[t]]$ verifies $g_0 = 0$, then $f(g(X)) = f \circ g(X)$ where $f \circ g \in \mathbb{K}_0[[t]]$ is defined by $f \circ g(t) = f(g(t))$. In particular, $(1 + X)^{-1} := \sum_{n \geq 0} (-1)^n X^n$, $\log(1 + X) := \sum_{n > 0} (-1)^{n+1} \frac{X^n}{n}$ and $\exp(X) := \sum_{n \geq 0} \frac{X^n}{n!}$ are defined. We shall need the following standard supplement to the functional calculus in subsection 2.3.

Lemma 1.1. *Let $\alpha \in F(M, M)$, $f(t) \in \mathbb{K}_0[[t]]$, and $X \in F(M)_0$ be such that $\delta(X) = 0$. Then $\zeta_\alpha(f(X)) = f'(X)\zeta_\alpha(X)$.*

Proof. We leave to the reader to check that it reduces to the case $f = t^n$ and to the fact that ζ_α is a derivation. $\square$

Lemma 1.2. *Suppose that N is a free $\mathbb{K}$ -module of finite rank (p, q) . Then $F(M, N)$ is a free $F(M)$ -module of finite rank (p, q) . Let $(\phi_i)_{i \in I}$ be a family of elements of $F(M, N)$. Then it is a basis of $F(M, N)$ if and only if the family $(\delta(\phi_i))_{i \in I}$ is a basis of N .*

Proof. The inclusion $N \subset F(M, N)$ induces an isomorphism $F(M) \otimes N \simeq F(M, N)$. Let $v_i := \delta(\phi_i)$. Obviously, if $(\phi_i)_{i \in I}$ is a basis of $F(M, N)$ then $(v_i)_{i \in I}$ is a basis of N . Conversely, suppose that $(v_i)_{i \in I}$ is a basis of N . By the inclusion $N \subset F(M, N)$ it is a basis of $F(M, N)$. Write $\phi_j = \sum v_i g_j^i$ with $g_j^i \in F(M)$. The square matrix $g := (g_j^i)$ is even. The matrix $\delta(g)$ is the identity I . By functional calculus g is invertible with inverse $(I + (g - I))^{-1} = I + (I - g) + (I - g)^2 + \dots$. Thus $(\phi_i)_{i \in I}$ is a basis of $F(M, N)$. $\square$

1.4. Berezinians and supertraces. In this subsection, we consider a $\mathbb{K}$ -module N which is free of finite rank.

Inside the commutative superalgebra $F(N, S(\Pi N))$ we consider the subalgebra $S(\Pi N) \otimes S(N^*) = S(\Pi N \oplus N^*)$, and we still denote by π the element of $S(\Pi N \oplus N^*)$ which corresponds to $\pi \in \text{Hom}(N, \Pi N)$. We have $\pi^2 = 0$. Following Manin [15, ch. 3.4], we define the *Berezinian of the module N*

$$(20) \quad \text{Ber}_{\mathbb{K}}(N) := \ker \pi / \text{im } \pi$$

as the homology of the multiplication by π in $S(\Pi N \oplus N^*)$. We write $\text{Ber } N$ or $\text{Ber}(N)$ for $\text{Ber}_{\mathbb{K}}(N)$ if $\mathbb{K}$ is clear from the context.

Let $(e_i)_{1 \leq i \leq n}$ be a totally ordered basis of N , and $(x^i)_{1 \leq i \leq n}$ be the dual basis of N^* . We have $\pi = \sum \pi e_i x^i$. Let $b_i = \pi e_i$ if e_i is even, and $b_i = x^i$ if e_i is odd. The element $b_1 \cdots b_n \in S^n(\Pi N \oplus N^*)$ is annihilated by π . We denote by

$$(21) \quad D(e_1, \dots, e_n) = b_1 \cdots b_n \mod \pi S(\Pi N \oplus N^*)$$

its class in $\text{Ber}(N)$. We recall the following fundamental fact from Manin [15, ch. 3.4.7]:

Lemma 1.3. *The $\mathbb{K}$ -module $\text{Ber}(N)$ is free of rank 1, with basis $D(e_1, \dots, e_n)$.*

The module $\text{Ber}(N)$ is a generalization of the top part of the exterior algebra in the free ungraded situation. Indeed, when $\mathbb{K} = \mathbb{K}_0$ and $N = N_0$ is free with a finite basis $(e_i)_{1 \leq i \leq n}$ consisting entirely of even elements, then $\text{Ber}(N)$ is the top degree part of the exterior algebra of N .

Remark 1.2. *Manin's definition of the Berezinian module is in fact $\Pi^p \text{Ber}(N)$, where (p, q) is the rank of N — see [15, ch. 3.4.7]. In our presentation, there is no advantage in doing this change of parity.*

If A is an associative algebra with unit, we denote by $A^\times$ the multiplicative group of invertible elements. We denote by $GL(N) = \text{End}(N)_{\mathbf{0}}^\times$ the group of even isomorphisms of N . The natural action of $g \in GL(N)$ in $\text{Ber}(N)$ is by an even invertible scalar $\text{Ber}(g) \in \mathbb{K}_{\mathbf{0}}^\times$ (or $\text{Ber}_N(g)$ if we need to specify) called the *Berezinian of g* .

Similarly, the natural Lie action of $X \in \text{End}(N)$ in $\text{Ber}(N)$ is denoted by $\text{str}(X)$ (or $\text{str}_N(X)$), and $\text{str} \in \text{End}(N)_{\mathbf{0}}^*$ is a 1-dimensional representation of $\text{End}(N)$ (considered as a Lie superalgebra) called the *supertrace*. In terms of a basis, we have (see [15, ch. 3.3])

$$(22) \quad \text{str}(X) = \sum (-1)^{p(e_i)(p(e_i)+p(X))} x^i (X e_i).$$

Berezinian modules have functorial properties which are obvious on the level of the defining complexes: $\text{Ber}(N) \simeq \text{Ber}((\Pi N)^*)$, $\text{Ber}(N \oplus N') \simeq \text{Ber}(N) \otimes \text{Ber}(N')$. Moreover, if $\mathbb{K}'$ is a commutative $\mathbb{K}$ -superalgebra, we have a canonical isomorphism

$$(23) \quad \text{Ber}_{\mathbb{K}'}(N \otimes_{\mathbb{K}} \mathbb{K}') \simeq \text{Ber}_{\mathbb{K}}(N) \otimes_{\mathbb{K}} \mathbb{K}'.$$

It is less obvious that there is a natural duality between $\text{Ber}(N^*)$ and $\text{Ber}(N)$:

Lemma 1.4. *There exists an unique $\mathbb{K}$ -bilinear map $\langle \cdot, \cdot \rangle: \text{Ber}(N^*) \times \text{Ber}(N) \rightarrow \mathbb{K}$ such that, for any basis $(e_i)_{1 \leq i \leq n}$ of N , denoting by $(x^i)_{1 \leq i \leq n}$ the dual basis of N^* , we have*

$$(24) \quad \langle D(x^n, \dots, x^1), D(e_1, \dots, e_n) \rangle = 1.$$

Remark 1.3. *Note that we reversed the order of the basis of N^* to avoid signs. The following computational proof of lemma 1.4 is not very informative. It would be interesting to replace it by a more categorical argument.*

Proof. The lemma is equivalent to the following formula:

$$(25) \quad \text{Ber}(g) = \text{Ber}(g^*) \text{ for } g \in GL(N),$$

where $g^* \in \text{End}(N^*)$ is the transpose of g . This identity is given in [15, ch. 3.3.6]. $\square$

In other terms, there is a canonical isomorphism

$$(26) \quad \text{Ber}(N^*) \otimes \text{Ber}(N) \rightarrow \mathbb{K}.$$

We denote by

$$(27) \quad \mathbf{d} := D(x^n, \dots, x^1) \otimes D(e_1, \dots, e_n) \in \text{Ber}(N^*) \otimes \text{Ber}(N)$$

the inverse image of 1, and we call it the *canonical element*.

Let M be a $\mathbb{K}$ -module. Since N is free of finite rank, the injection $N \subset F(M, N)$ induces isomorphisms $F(M) \otimes N \simeq F(M, N)$ and $F(M, \text{End}_{\mathbb{K}}(N)) \simeq \text{End}_{F(M)}(F(M, N))$. Let $X \in F(M, \text{End}_{\mathbb{K}}(N))_{\mathbf{0}}$ be such that $\delta(X) = 0$. We have (see [15, ch. 3.3] for a similar statement):

$$(28) \quad \text{Ber}_{F(M, N)}(\exp(X)) = \exp(\text{str}_{F(M, N)}(X)).$$

Note that the two sides of this equation are well defined elements of $F(M)$. For the left hand side, this is because $\exp(X) \in GL(F(M, N))$. For the right hand side, this is because $\delta(\text{str}(X)) = 0$.

Consider in addition $r = \sum_{n \geq 0} r_n t^n \in \mathbb{K}_{\mathbf{0}}[[t]]$ such that $r_0 = 1$. Since $\delta(r(X)) = 1$, the element $r(X)$ is invertible and has a Berezinian $\text{Ber}(r(X)) \in F(M)$. Define $w = \log(r) \in t\mathbb{K}_{\mathbf{0}}[[t]]$ by the formula:

$$(29) \quad w = \log(1 + (r - 1)) = (r - 1) - \frac{(r - 1)^2}{2} + \dots = r_1 t + (r_2 - \frac{r_1^2}{2}) t^2 + \dots$$

By functional calculus, we have $r(X) = \exp(w(X))$ and, using (28), we find

$$(30) \quad \text{Ber}(r(X)) = \exp(\text{str}(w(X))).$$

1.5. Dualizing modules. Let $y^1, \dots, y^n$ be formal even or odd variables. We denote by $\mathbb{K}[[y^1, \dots, y^n]]$ the superalgebra of formal series, that is the commutative superalgebra of formal sums with coefficients in $\mathbb{K}$ of monomials $(y^1)^{m_1} \dots (y^n)^{m_n}$, with $m_i \in \mathbb{N}$, and $m_i \leq 1$ if y^i is odd. If p is the number of even variables and q the number of odd variables, we say that the dimension of $\mathbb{K}[[y^1, \dots, y^n]]$ is (p, q) . We denote by $\delta \in \text{Hom}(\mathbb{K}[[y^1, \dots, y^n]], \mathbb{K})$ the algebra homomorphism which is zero on the y^i .

Remark 1.4. *Let $\mathbb{K}$ be a field. Then the algebra $\mathbb{K}[[y^1, \dots, y^n]]$ is local, and δ is its unique algebra homomorphism with values in $\mathbb{K}$. As we prefer not to make this assumption on $\mathbb{K}$, we specify δ and consider $\mathbb{K}[[y^1, \dots, y^n]]$ as an augmented algebra.*

Augmented algebras. Recall that an associative $\mathbb{K}$ -superalgebra A with unit is called *augmented* if it is provided with a $\mathbb{K}$ -algebra morphism $\delta : A \rightarrow \mathbb{K}$, the augmentation. We use the notation $A^+ = \ker \delta \subset A$. The map δ induces an isomorphism $A/A^+ \simeq \mathbb{K}$ and provides $\mathbb{K}$ with an A -algebra structure.

We provide A with the A^+ -adic topology: a linear endomorphism $X \in \text{End}(A)$ is continuous if for all $n \in \mathbb{N}$, there exists $m \in \mathbb{N}$ such that $X((A^+)^m) \subset (A^+)^n$. The A -module of continuous derivations of A is denoted by $\mathcal{T}(A)$. A linear form $w \in A^*$ is continuous if there exists $n \in \mathbb{N}$ such that $w|_{(A^+)^n} = 0$. We denote by $A_* \subset A^*$ the continuous dual (it is also called the space of *distributions with support* $\{\delta\}$).

If $\mathcal{V}$ is an A -module, we consider the $\mathbb{K}$ -module $V := \mathcal{V}/A^+\mathcal{V} = \mathcal{V} \otimes_A \mathbb{K}$, and we denote by $\bar{\delta} = \text{id}_{\mathcal{V}} \otimes_A \delta$ the quotient map $\mathcal{V} \rightarrow V$. We denote by $\mathfrak{a}$ the $\mathbb{K}$ -module of δ -derivations $A \rightarrow \mathbb{K}$.

Example 1.1. (Basic example). *Let $A = \mathbb{K}[[y^1, \dots, y^n]]$. We use the notation $\frac{\partial}{\partial y^i}$ for the continuous derivation of $\mathbb{K}[[y^1, \dots, y^n]]$ such that $\frac{\partial y^j}{\partial y^i} = \delta_i^j$. These derivation form a basis of the $\mathbb{K}[[y^1, \dots, y^n]]$ -module $\mathcal{T}(\mathbb{K}[[y^1, \dots, y^n]])$. The elements $e_i := (-1)^{p(y^i)} \delta \circ \frac{\partial}{\partial y^i}$ provide a basis of $\mathfrak{a}$. The images of y^i in $A^+/(A^+)^2$ form the dual basis of $\mathfrak{a}^*$, and the module $\bigoplus_{i=1}^n \mathbb{K}y^i$ is a supplementary factor for $(A^+)^2$ in A^+ .*

Definition 1.1. *We say that an augmented $\mathbb{K}$ -superalgebra A is formally smooth of dimension (p, q) if it is isomorphic to an augmented algebra $\mathbb{K}[[y^1, \dots, y^n]]$ of dimension (p, q) .*

Thus, if A is formally smooth, the choice of an isomorphism $\theta^* : A \rightarrow \mathbb{K}[[y^1, \dots, y^n]]$ is equivalent to the choice of the corresponding elements $x^i := \theta^{*-1}(y^i) \in A^+$. We write $A = \mathbb{K}[[x^1, \dots, x^n]]$, and call the family $(x^i)_{1 \leq i \leq n}$ “a system of local coordinates at the point $\{\delta\}$ ”.

The dualizing module of a formally smooth augmented algebra. For the rest of this section, we consider a formally smooth augmented $\mathbb{K}$ -superalgebra A of dimension (p, q) . Thus $\mathfrak{a}$ is free of rank (p, q) over $\mathbb{K}$ and $\mathcal{T}(A)$ is free of rank (p, q) over A . The space A_* is a cocommutative coalgebra with counit δ , and A can be identified to the dual of A_* .

The dual $\text{Hom}_A(\mathcal{T}(A), A)$ is denoted by $\Omega_{ev}^1(A)$: it is the space of *even 1-differential forms*. We denote by $\Omega^1(A)$ the space $\Pi \Omega_{ev}^1$: it is the space of *1-differential forms*. For $f \in A$, we denote by $d_{ev}f$ the element of $\Omega_{ev}^1(A)$ such that $\mathcal{T}(A) \ni \zeta \mapsto \pm \zeta(f)$, and $df := \pi d_{ev}f \in \Omega^1(A)$.

Following [25], we define the *dualizing module* $K(A)$ by the formula

$$(31) \quad K(A) = \text{Ber}_A(\Omega_{ev}^1(A)).$$

It is a free A -module of rank 1.

Let $(x^i)_{1 \leq i \leq n}$ be a choice of local coordinates at the point $\{\delta\}$. The family $(d_{ev}x^i)$ is a basis of $\Omega_{ev}^1(A)$ with dual basis of $\mathcal{T}(A)$ the vector fields $\frac{\partial}{\partial x^i}$. Recall that — see formula (21) — the class of $b_1 \dots b_n$, with $b_i = dx^i$ if x^i is even, or $b_i = \frac{\partial}{\partial x^i}$ if x^i is odd, is a basis of $K(A)$ which is denoted by $D(d_{ev}x^1, \dots, d_{ev}x^n)$.

Thus, the $\mathbb{K}$ -module $K(A)/A^+K(A)$ is free of rank 1.

Remark 1.5. Since $\bar{\delta} : \Omega_{ev}^1(A)/A^+\Omega_{ev}^1(A) \rightarrow \mathfrak{a}^*$ is an isomorphism, we obtain from (23) an isomorphism

$$(32) \quad \bar{\delta} : K(A)/A^+K(A) \rightarrow \text{Ber}_{\mathbb{K}}(\mathfrak{a}^*).$$

In coordinates, for $\phi \in A$, we have $\bar{\delta}(D(d_{ev}x^1, \dots, d_{ev}x^n)\phi) = D(x^1, \dots, x^n)\delta(\phi)$.

Tensorial modules and divergence. Recall that $\mathcal{T}(A)$ is a Lie $\mathbb{K}$ -subalgebra of $\text{End}_{\mathbb{K}}(A)$.

Definition 1.2. We say that an A -module $\mathcal{V}$ endowed with a $\mathcal{T}(A)$ -module structure is a $\mathcal{T}(A)$ - A -module if we have

$$(33) \quad \zeta(fv) = \zeta(f)v \pm f\zeta(v) \text{ for all } f \in A, v \in \mathcal{V}, \zeta \in \mathcal{T}(A).$$

The action of ζ is called the Lie derivative with respect to ζ in $\mathcal{V}$, and is traditionally denoted by $\mathcal{L}(\zeta)$.

If $\mathcal{V}$ is a $\mathcal{T}(A)$ - A -module, so are the module $\Pi\mathcal{V}$ and the dual $\mathcal{V}^* = \text{Hom}_A(\mathcal{V}, A)$, each one with the natural actions. If $\mathcal{V}$ and $\mathcal{W}$ are $\mathcal{T}(A)$ - A -modules, so are $\mathcal{V} \otimes_A \mathcal{W}$ and $\text{Hom}_A(\mathcal{V}, \mathcal{W})$, with the natural actions.

For instance, A with the Lie derivative $\mathcal{L}(\zeta) = \zeta$ is a $\mathcal{T}(A)$ - A -module.

Example 1.2. Another important example for this text is $\mathcal{T}(A)$, with the Lie derivative $\mathcal{L}(\zeta) = \text{ad}(\zeta)$.

It follows that the tensor spaces, obtained from $\Omega_{ev}^1(A)$, $\Omega^1(A) \equiv \Pi\Omega_{ev}^1(A)$, $\mathcal{T}(A)$ and $\Pi\mathcal{T}(A)$ by taking tensor products over A and $\mathcal{T}(A)$ -invariant subquotients, are naturally $\mathcal{T}(A)$ - A -modules. We call these $\mathcal{T}(A)$ - A -modules *tensorial modules*.

Remark 1.6. In particular, since $\pi \in \text{Hom}_A(\Omega_{ev}^1(A), \Omega^1(A))$ is annihilated by the Lie derivatives, $K(A)$ inherits a Lie derivative and is a tensorial module.

We shall need the following relation (see [12] for a similar statement):

$$(34) \quad \mathcal{L}(f\zeta)(D) = f\mathcal{L}(\zeta)(D) \pm \zeta(f)D, \text{ for } f \in A, \zeta \in \mathcal{T}(A), D \in K(A).$$

Let $D \in K(A)$ be a basis. The *divergence* $\text{div}_D(\zeta) \in A$ of $\zeta \in \mathcal{T}(A)$ with respect to the basis D is defined by

$$(35) \quad \mathcal{L}(\zeta)(D) = \text{div}_D(\zeta)D.$$

Invariance by translation. Let M be a free $\mathbb{K}$ -module of finite rank (p, q) . We specialize to the case $A = F(M)$. Let $(e_i)_{1 \leq i \leq n}$ be a totally ordered basis of M , and $(x^i)_{1 \leq i \leq n}$ the corresponding dual basis of M^* . This provides an identification $F(M) = \mathbb{K}[[x^1, \dots, x^n]]$, and $F(M)$ is formally smooth with local coordinates $(x^i)_{1 \leq i \leq n}$.

We say that an element $d \in K(F(M))$ is *invariant by translations* if $\mathcal{L}(\zeta_m)d = 0$ for all $m \in M$ (ζ_m is the derivative in the direction m defined in (15)).

Lemma 1.5. The $\mathbb{K}$ -submodule of $K(F(M))$ of elements which are invariant by translations is free of rank 1. A basis of this module is also a basis of $K(F(M))$ considered as an $F(M)$ -module.

Proof. In coordinates, $D(d_{ev}x^1, \dots, d_{ev}x^n)$ is such a basis. $\square$

For a basis d invariant by translation, we write simply div for div_d . Let $\sum \zeta^i \frac{\partial}{\partial x^i}$ be a derivation of $\mathbb{K}[[x^1, \dots, x^n]]$. The following well known formula follows from invariance by translations and formula (34):

$$(36) \quad \text{div}\left(\sum \zeta^i \frac{\partial}{\partial x^i}\right) = \sum \pm \frac{\partial \zeta^i}{\partial x^i}.$$

Isomorphisms of augmented algebras. Let A and B be two formally smooth augmented superalgebras and $\theta^* : B \rightarrow A$ an isomorphism of augmented algebras. By transposition,

θ^* induces an isomorphism of Lie $\mathbb{K}$ -superalgebras $\theta_* : \mathcal{T}(A) \rightarrow \mathcal{T}(B)$, and of $\mathbb{K}$ -modules $\theta^* : \Omega_{ev}^1(B) \rightarrow \Omega_{ev}^1(A)$. We denote also by $\theta^* : \mathcal{T}(B) \rightarrow \mathcal{T}(A)$ the inverse of the map θ_* . This induces $\mathbb{K}$ -isomorphisms θ^* between various tensor spaces related to B and A , and in particular a bijection

$$(37) \quad \theta^* : K(B) \rightarrow K(A)$$

which satisfy the relations

$$(38) \quad \theta^*(f\omega) = \theta^*(f)\theta^*(\omega) \text{ for } f \in B \text{ and } \omega \in K(B),$$

$$(39) \quad \theta^*(\zeta\omega) = \theta^*(\zeta)\theta^*(\omega) \text{ for } \zeta \in \mathcal{T}(B) \text{ and } \omega \in K(B).$$

1.6. Dualizing modules and duality. Let A be a formally smooth augmented superalgebra. Among the many possible tensorial modules, the dualizing module $K(A)$ is specially important because of its role in integration and duality. This is explained in the case of supermanifolds (algebraic, complex analytic, smooth) in [15, 24, 25]. In our formal context, see the work of Chemla [5–7]. To study Gorelik elements, we need only the simple particular case which we describe below.

Assume that A has dimension $(0, q)$. Let $x^1, \dots, x^q$ be odd coordinates. We have $A = \mathbb{K}[x^1, \dots, x^q]$ —in this case formal series are polynomials. As a $\mathbb{K}$ -module, A is free of finite rank 2^q . It implies that the dual A^* is equal to the continuous dual A_* . We have $K(A) = S^q(\mathcal{T}(A))$. Recall that the vector fields $\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^q}$ are a basis of $\mathcal{T}(A)$. Any element of $K(A)$ can be written in an unique manner as $\frac{\partial}{\partial x^1} \cdots \frac{\partial}{\partial x^q} \phi$ with $\phi \in A$. Berezin integral $\int : K(A) \rightarrow \mathbb{K}$ is defined by the formula

$$(40) \quad \int \frac{\partial}{\partial x^1} \cdots \frac{\partial}{\partial x^q} \phi = \delta \left(\frac{\partial}{\partial x^1} \cdots \frac{\partial}{\partial x^q} (\phi) \right).$$

The main point, due to Berezin, is that Berezin integral does not depend on the choice of the coordinates $x^1, \dots, x^q$. Berezin integral provides an isomorphism of $\mathcal{T}(A)$ – A -modules $\psi_A : K(A) \rightarrow A_*$ by the formula

$$(41) \quad \psi_A(\omega)(\phi) = \int \omega \phi \quad \text{for } \omega \in K(A), \phi \in A.$$

The choice of coordinates identifies A and $S(\mathfrak{a})^*$, and A_* and $S(\mathfrak{a})$. We write ψ_A using this identification, and the elements $e_i \in \mathfrak{a}$ such that $x^j(e_i) = \delta_i^j$ — note that $e_i = -\delta \circ \frac{\partial}{\partial x^i}$.

Lemma 1.6. *We have*

$$\psi_A \left(\frac{\partial}{\partial x^1} \cdots \frac{\partial}{\partial x^q} \phi \right) = (-1)^q e_1 \dots e_q \phi.$$

Proof. Since ψ_A is a morphism of A modules, it is sufficient to prove the lemma for $\phi = 1$. We have $\psi_A(\frac{\partial}{\partial x^1} \cdots \frac{\partial}{\partial x^q})(\phi') = 0$ if $\phi' \in \mathbb{K}[x^1, \dots, x^q]$ is of degree smaller than q , and $\psi_A(\frac{\partial}{\partial x^1} \cdots \frac{\partial}{\partial x^q})(x^q \cdots x^1) = 1$. This proves the lemma. $\square$

Let B another augmented algebra and $\theta^* : B \rightarrow A$ and isomorphism of augmented algebras. It is the transpose of a coalgebra isomorphism $\theta : A_* \rightarrow B_*$. Recall from (37) the isomorphism $\theta^* : K(B) \rightarrow K(A)$.

Lemma 1.7. *We have $\psi_B = \theta \circ \psi_A \circ \theta^*$.*

Proof. For $\omega \in K(B)$, we have $\int \theta^*(\omega) = \int \omega$. The lemma follows from (38). $\square$

2. DUALIZING MODULES ON FORMAL LIE SUPERGROUPS.

Let $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ be a Lie $\mathbb{K}$ -superalgebra, and $U(\mathfrak{g})$ be its enveloping algebra. We denote by $j : \mathfrak{g} \rightarrow U(\mathfrak{g})$ the canonical map. Since $\mathbb{K}_0$ contains $\mathbb{Q}$, j is injective (cf. [8, 18]).

2.1. Enveloping algebras as coalgebras. We use the notation $U^+(\mathfrak{g})$ for the ideal generated by $\mathfrak{g}$.

Recall that $U(\mathfrak{g})$ is a cocommutative Hopf superalgebra. For $u \in U(\mathfrak{g})$, we write

$$(42) \quad \Delta_{U(\mathfrak{g})}(u) = \sum u_i \otimes u'_i$$

for its coproduct, and we denote by $S_{U(\mathfrak{g})} : U(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ the antipode. We use the notation $F(G)$ for the commutative superalgebra $U(\mathfrak{g})^*$ (dual of the commutative coalgebra $U(\mathfrak{g})$) provided with the augmentation $\delta : f \rightarrow f(1)$. This can be considered as the *definition of a formal Lie supergroup* G : $F(G)$ is the space of formal functions defined in a formal neighborhood of $1 \in G$, and δ is the evaluation at 1.

For a $\mathbb{K}$ -module N , we define in a similar way $F(G, N) = \text{Hom}(U(\mathfrak{g}), N)$, considered as the N -valued functions on G . As in subsection 1.2, $F(G, N)$ is an $F(G)$ -module, and, if $\mu : N_1 \otimes N_2 \rightarrow N$ is a map of $\mathbb{K}$ -modules, there is a corresponding map $\tilde{\mu} : F(G, N_1) \otimes_{F(G)} F(G, N_2) \rightarrow F(G, N)$. In particular, if N is a K -superalgebra, $F(G, N)$ is an $F(G)$ -superalgebra.

If f_1 and f_2 are in $\text{End}(U(\mathfrak{g}))$ we use the notation $f_1 * f_2$ for the product in $F(G, U(\mathfrak{g})) = \text{End}(U(\mathfrak{g}))$. Formulas (14) generalize in two versions, one for *left* and one for *right action*. Here are the details. Let $\phi \in \text{End}(U(\mathfrak{g}))$. We use the notation $c_\phi^\lambda = \phi * \text{id}_{U(\mathfrak{g})}$ and we denote by ζ_ϕ^λ the transpose map acting in $F(G, N)$. For $u \in U(\mathfrak{g})$ and $g \in F(G, N)$, we have

$$(43) \quad c_\phi^\lambda(u) = \sum \phi(u_i)u'_i \text{ and } \zeta_\phi^\lambda(g) = \pm g \circ c_\phi^\lambda.$$

Similarly, we use the notation $c_\phi^\rho = \text{id}_{U(\mathfrak{g})} * \phi$ and we denote by ζ_ϕ^ρ the transpose map acting in $F(G, N)$. Thus we have

$$(44) \quad c_\phi^\rho(u) = \sum \pm u_i \phi(u'_i) \text{ and } \zeta_\phi^\rho(g) = \pm g \circ c_\phi^\rho.$$

Note that $\phi = S_{U(\mathfrak{g})} * c_\phi^\rho = c_\phi^\lambda * S_{U(\mathfrak{g})}$, and that $c_\phi^\rho = \text{id}_{U(\mathfrak{g})} * c_\phi^\lambda * S_{U(\mathfrak{g})}$.

Let $v \in U(\mathfrak{g})$. We denote by l_v the left multiplication by v in $U(\mathfrak{g})$. Identifying v with the constant function on G with value v we get $c_v^\lambda = l_v$. Similarly, $c_v^\rho(u) = \pm uv$, ζ_v^λ is the right invariant differential operator whose evaluation at 1 is equal to v , and ζ_v^ρ is the left invariant differential operator whose evaluation at 1 is equal to v . For $a \in \mathfrak{g}$, we will simply write $l_a := l_{j(a)}$, $\zeta_a^\lambda := \zeta_{j(a)}^\lambda$ and $\zeta_a^\rho := \zeta_{j(a)}^\rho$.

Let $f \in F(G)$. Identifying it with a scalar valued element of $\text{End}(U(\mathfrak{g}))$, we get $c_f^\rho = c_f^\lambda$. This provides $U(\mathfrak{g})$ with a structure of $F(G)$ -module. Accordingly, for $u \in U(\mathfrak{g})$, we also write fu for $c_f(u)$.

An element $\alpha \in F(G, \mathfrak{g})$ is called a *vector field on G*.

Remark 2.1. Both ζ_α^λ and ζ_α^ρ are elements of $\mathcal{T}(F(G))$.

Let $\beta \in F(G, \mathfrak{g})$ be a vector field. We have

$$(45) \quad [\zeta_\alpha^\rho, \zeta_\beta^\rho] = \zeta_\gamma^\rho \text{ with } \gamma = \zeta_\alpha^\rho(\beta) - \pm \zeta_\beta^\rho(\alpha) + [\alpha, \beta],$$

$$(46) \quad [\zeta_\alpha^\lambda, \zeta_\beta^\lambda] = \zeta_{\gamma'}^\lambda \text{ with } \gamma' = \zeta_\alpha^\lambda(\beta) - \pm \zeta_\beta^\lambda(\alpha) - [\alpha, \beta],$$

$$(47) \quad [\zeta_\alpha^\lambda, \zeta_\beta^\rho] = \zeta_{\gamma_1}^\rho - \pm \zeta_{\gamma_2}^\lambda, \text{ with } \gamma_1 = \zeta_\alpha^\lambda(\beta) \text{ and } \gamma_2 = \zeta_\beta^\rho(\alpha).$$

These formulas include as particular cases formulas (16) when $\mathfrak{g}$ is commutative, and the formulas $[\zeta_a^\rho, \zeta_b^\rho] = \zeta_{[a,b]}^\rho$, $[\zeta_a^\lambda, \zeta_b^\lambda] = -\zeta_{[a,b]}^\lambda$ and $[\zeta_a^\lambda, \zeta_b^\rho] = 0$ for $a \in \mathfrak{g}$ and $b \in \mathfrak{g}$. We have similar relations for the coderivations c_α^ρ and c_α^λ of $U(\mathfrak{g})$. We note the following classical proposition.

Proposition 2.1. *The map $\alpha \rightarrow c_\alpha^\rho$ is an isomorphism from $F(G, \mathfrak{g})$ to the space of coderivations of $U(\mathfrak{g})$.*

Proof. Since $\mathbb{K}$ contains $\mathbb{Q}$, $\mathfrak{g}$ is the subspace of primitive elements of $U(\mathfrak{g})$. As in [20, theorem 1], this implies the proposition. $\square$

2.2. The exponential map of a formal Lie supergroup. Recall from [8], [18] that the symmetrization $\beta : S(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ is a coalgebra isomorphism such that $\beta(1) = 1$ and $\beta(a) = j(a)$ for $a \in \mathfrak{g}$.

Remark 2.2. *The transpose map $\beta^* : F(G) \rightarrow F(\mathfrak{g})$ is an isomorphism of augmented algebras.*

We denote also by $\beta^* : F(G, N) \rightarrow F(\mathfrak{g}, N)$ the map defined by $\beta^*(g) = g \circ \beta$ for $g \in F(G, N)$.

We consider β as a formal diffeomorphism from a neighborhood of $0 \in \mathfrak{g}$ to a neighborhood of $1 \in G$. This is the formal analog of the usual exponential map. It can also be written as an exponential in our formal setting: extend j to an element of the algebra $F(\mathfrak{g}, U(\mathfrak{g}))$ by assigning the value zero on $S^n(\mathfrak{g})$ for $n \neq 1$. The sum $\exp(j) = \sum_{n \geq 0} \frac{j^n}{n!}$ is locally finite and $\beta = \exp(j)$.

2.3. The generic point of a Lie superalgebra. In this subsection, we collect some results on $F(\mathfrak{g})$ needed later. We write simply $x = x_{\mathfrak{g}} \in F(\mathfrak{g}, \mathfrak{g})$ the generic point of $\mathfrak{g}$.

We consider $\text{ad } x \in \text{End}_{F(\mathfrak{g})}(F(\mathfrak{g}, \mathfrak{g}))$. For $p = \sum_{n \geq 0} p_n t^n \in \mathbb{K}_{\mathbf{0}}[[t]]$, we can define $p(\text{ad } x) \in \text{End}_{F(\mathfrak{g})}(F(\mathfrak{g}, \mathfrak{g}))$ by a locally finite sum as in (19). For any vector field $\alpha \in F(\mathfrak{g}, \mathfrak{g})$, we obtain a new vector field $p(\text{ad } x)(\alpha)$. As in [18], the vector fields

$$(48) \quad p(\text{ad } x)(a) = \sum_{n \geq 0} p_n \text{ad}^n(x)(a) \text{ with } a \in \mathfrak{g},$$

their corresponding coderivations $c_{p(\text{ad } x)(a)}$ of $S(\mathfrak{g})$, and the transpose operators $\zeta_{p(\text{ad } x)(a)}$ of $F(\mathfrak{g}, N)$ (N is a $\mathbb{K}$ -module), play a special role. We recall how to compute $p(\text{ad } x)(a)$. Let $b_1, b_2, \dots, b_n$ in $\mathfrak{g}$. We have (see [18])

$$(49) \quad p(\text{ad } x)(a) : b_1 \cdots b_n \mapsto p_n \sum_{s \in S_n} \pm \text{ad } b_{s(1)} \cdots \text{ad } b_{s(n)}(a).$$

In the rest of this section, we suppose that $\mathfrak{g}$ is free of finite rank. The Lie superalgebra $F(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g} \otimes F(\mathfrak{g})$ is free of finite rank over $F(\mathfrak{g})$. Thus, the divergence $\text{div}(\zeta_{p(\text{ad } x)a}) \in F(\mathfrak{g})$ of the derivation associated to the vector field $p(\text{ad } x)(a)$ is defined in (36). Similarly, for $w \in \mathbb{K}_{\mathbf{0}}[[t]]$, the function $\text{str}(w(\text{ad } x)) \in F(\mathfrak{g})$ is defined. We state some results relating these objects.

Proposition 2.2.

$$(50) \quad \text{div}(\zeta_{p(\text{ad } x)a}) = -\text{str} \left(\frac{p(\text{ad } x) - p(0)}{\text{ad } x} \text{ad } a \right).$$

Proof. It is sufficient to prove it for $p(t) = t^n$. We pick a basis $(e_i)_{i \in I}$ of $\mathfrak{g}$ and the corresponding dual basis $(x^i)_{i \in I}$ of $\mathfrak{g}^*$. Let ζ_{e_i} the derivative in the direction e_i . We have

$$(51) \quad \begin{aligned} \zeta_{e_i}(\text{ad}^n(x)(a)) &= \sum_{p+q=n-1} \text{ad}^p(x) \text{ad}(e_i) \text{ad}^q(x)(a) \\ &= \sum_{p+q=n-1} \text{ad}^p(x) ([e_i, \text{ad}^q(x)(a)]) \\ &= - \sum_{p+q=n-1} \text{ad}^p(x) \text{ad}(\text{ad}^q(x)(a))(e_i). \end{aligned}$$

Applying formula (36), we get

$$(52) \quad \text{div}(\zeta_{\text{ad}^n(x)(a)}) = -\text{str} \left(\sum_{p+q=n-1} \text{ad}^p(x) \text{ad}(\text{ad}^q(x)(a)) \right).$$

Using the notations $X := \text{ad } x$, $A := \text{ad } a$ and the relation $\text{ad}(\text{ad}^n(x)(a)) = \text{ad}^n(X)(A)$, we obtain

$$(53) \quad \text{div}(\zeta_{\text{ad}^n(x)(a)}) = -\text{str} \left(\sum_{p+q=n-1} X^p \text{ad}^q(X)(A) \right).$$

Let l and r be the left and right multiplication by X in $\text{End}_{F(\mathfrak{g})}(F(\mathfrak{g}, \mathfrak{g}))$. Since $\text{ad } X = l - r$, we obtain

$$(54) \quad \text{div}(\zeta_{\text{ad}^n(x)(a)}) = -\text{str} \left(\sum_{p+q=n-1} l^p (l - r)^q(A) \right).$$

For all $Y \in \text{End}_{F(\mathfrak{g})}(F(\mathfrak{g}, \mathfrak{g}))$ we have $\text{str}((l - r)Y) = \text{str}(XY - YX) = 0$. We obtain

$$(55) \quad \text{div}(\zeta_{\text{ad}^n(x)(a)}) = -\text{str}(l^{n-1}(A)) = -\text{str}(\text{ad}^{n-1}(x) \text{ad } a),$$

which is what we want. $\square$

Proposition 2.3.

$$(56) \quad \zeta_{p(\text{ad } x)(a)}(\text{str}(w(\text{ad } x))) = \text{str}(p(0)w'(\text{ad } x) \text{ad } a).$$

Proof. It is sufficient to prove it for $w = t^n$. Using the same notations and arguments in the proof of proposition 2.2 we have

$$(57) \quad \zeta_{p(\text{ad } x)(a)}(\text{str}(\text{ad}^n(x))) = \sum_{p+q=n-1} \text{str}(\text{ad}^p(x) \text{ad}(p(\text{ad } x)(a)) \text{ad}^q(x))$$

$$(58) \quad = \sum_{p+q=n-1} \text{str}(X^p p(\text{ad } X)(A) X^q)$$

$$(59) \quad = \sum_{p+q=n-1} \text{str}(l^p r^q p(l - r)(A)).$$

Since $p(l - r) - p(0)$ is divisible by $l - r$, we obtain

$$(60) \quad \zeta_{p(\text{ad } x)(a)}(\text{str}(\text{ad}^n x)) = \sum_{p+q=n-1} \text{str}(p(0)l^p r^q(A))$$

$$(61) \quad = \sum_{p+q=n-1} \text{str}(p(0)X^{p+q}A) = \text{str}(p(0)nX^{n-1}A),$$

which is what we want. $\square$

Let $r(t) = 1 + r_1 t + \dots \in \mathbb{K}_0[[t]]$. Since $r(\text{ad } x)$ is an even invertible endomorphism of $\mathfrak{g} \otimes F(\mathfrak{g})$, the Berezinian $\text{Ber } r(\text{ad } x) \in F(\mathfrak{g})$ is defined. Recall from (29) the definition of $w = \log(r)$. From (30) we obtain

$$(62) \quad \text{Ber } r(\text{ad } x) = \exp(\text{str}(w(\text{ad } x))).$$

Proposition 2.4.

$$(63) \quad \zeta_{p(\text{ad } x)(a)}(\text{Ber } r(\text{ad } x)) = \text{str}(p(0)w'(\text{ad } x) \text{ad } a) \text{Ber } r(\text{ad } x).$$

Proof. By lemma 1.1 applied with $f(t) = \exp(t)$ and $X = \text{str}(w(\text{ad } x))$, we have

$$(64) \quad \zeta_{p(\text{ad } x)(a)}(\text{Ber } r(\text{ad } x)) = \zeta_{p(\text{ad } x)(a)}(\text{str}(w(\text{ad } x))) \text{Ber } r(\text{ad } x).$$

Then the result follows from proposition 2.3. $\square$

2.4. The Jacobian of the exponential map. In this subsection we suppose that $\mathfrak{g}$ is free of finite rank as a $\mathbb{K}$ -module. The algebra $F(G)$ is formally smooth, and the dualizing module $K(F(G))$ is a free $F(G)$ -module of rank 1.

Remark 2.3. Let $\mathcal{V}$ be a $\mathcal{T}(F(G)) - F(G)$ -module (see definition 1.2). By composing the Lie derivative with the representations $a \rightarrow -\zeta_a^\lambda$ and $a \rightarrow \zeta_a^\rho$ of $\mathfrak{g}$, we get two commuting representations of $\mathfrak{g}$ in $\mathcal{V}$ (recall remark 2.1). They are called respectively the left and right actions of $\mathfrak{g}$.

Proposition 2.5. The map $F(G, \mathfrak{g}) \ni \alpha \rightarrow \zeta_\alpha^\rho \in \mathcal{T}(F(G))$ is an isomorphism of $F(G)$ -modules. It intertwines the left actions $-\zeta_a^\lambda$ and $\mathcal{L}(-\zeta_a^\lambda)$ of $a \in \mathfrak{g}$.

Proof. The first assertion follows from proposition 2.1. By definition of the Lie derivative in $\mathcal{T}(F(G))$ we have $\mathcal{L}(\zeta_a^\lambda)(\zeta_\alpha^\rho) = [\zeta_a^\lambda, \zeta_\alpha^\rho]$. Since $\zeta_\alpha^\rho(a) = 0$, it follows from formula (47) that we have $[\zeta_a^\lambda, \zeta_\alpha^\rho] = \zeta_\gamma^\rho$ with $\gamma = \zeta_a^\rho(\alpha)$. $\square$

Remark 2.4. By duality, we define an isomorphism $F(G, \mathfrak{g}^*) \ni \mu \mapsto \omega_{ev, \rho}^\mu \in \Omega_{ev}^1(F(G))$ by $\omega_{ev, \rho}^\mu(\zeta_\alpha^\rho) = \mu(\alpha)$. It intertwines the left actions $-\zeta_a^\lambda$ and $\mathcal{L}(-\zeta_a^\lambda)$ of $a \in \mathfrak{g}$. In particular, it defines a $\mathbb{K}$ -isomorphism $\mathfrak{g}^* \ni f \rightarrow \omega_{ev, \rho}^f$ into the space of left invariant differential forms.

Let $(e_i)_{1 \leq i \leq n}$ be a totally ordered basis of $\mathfrak{g}$, and $(x^i)_{1 \leq i \leq n}$ its dual basis. The derivations $\zeta_i^\rho := \zeta_{e_i}^\rho$ form a basis of $\mathcal{T}(F(G))$. Let $\omega_{ev, \rho}^i \in \Omega_{ev}^1(F(G))$ be the elements of the dual basis. We denote by $D^\rho = D(\omega_{ev, \rho}^1, \dots, \omega_{ev, \rho}^n)$ the corresponding basis of $K(F(G))$. It is invariant by the left action of $\mathfrak{g}$ in $K(F(G))$, and (recall (32)) $\bar{\delta}D^\rho = D(x^1, \dots, x^n) \in \text{Ber}(\mathfrak{g}^*)$. We obtain the following result.

Proposition 2.6. The $\mathbb{K}$ -module of elements of $K(F(G))$ invariant by the left action of $\mathfrak{g}$ is free of rank 1. A basis of this module is also a basis of $K(F(G))$ as an $F(G)$ -module.

Similarly, $d := D(d_{ev}x^1, \dots, d_{ev}x^n)$ is a basis of $K(F(\mathfrak{g}))$ invariant by translations (see lemma 1.5), and $\bar{\delta}d = D(x^1, \dots, x^n) \in \text{Ber}(\mathfrak{g}^*)$. The symmetrization β induces an isomorphism $\beta^* : K(F(G)) \rightarrow K(F(\mathfrak{g}))$, and the element $\beta^*(D^\rho)$ is a basis of $K(F(\mathfrak{g}))$. There exists a unique $J^\rho \in F(\mathfrak{g})_0$

$$(65) \quad \beta^*(D^\rho) = J^\rho d.$$

Since J^ρ does not depend on the choice of the basis $(e_i)_{1 \leq i \leq n}$, there is no ambiguity in the definition of J^ρ . The function J^ρ is called the *Jacobian of the exponential map in the left invariant frame*. The following formula is well known in the case of Lie groups.

Theorem 2.1. Let x be the generic point of $\mathfrak{g}$. We have

$$(66) \quad J^\rho = \text{Ber} \left(\frac{1 - e^{-\text{ad } x}}{\text{ad } x} \right).$$

Proof. We give two proofs. Both use the notations $r(t) = \frac{1-e^{-t}}{t}$, $p(t) = \frac{t}{e^t-1}$ and $w(t) = \log(r(t))$.

First proof. As in remark 2.4, replacing G with $\mathfrak{g}$, we identify $\Omega_{ev}^1(F(\mathfrak{g}))$ with $F(\mathfrak{g}, \mathfrak{g}^*)$. We recall the formula given in [18] for $\beta^*(\omega_{ev, \rho}^i) \in F(\mathfrak{g}, \mathfrak{g}^*)$. We consider $r(\text{ad } x) \in F(\mathfrak{g}, \text{End}(\mathfrak{g})) = \text{End}_{F(\mathfrak{g})}(F(\mathfrak{g}, \mathfrak{g}))$ as an element of $\text{GL}(F(\mathfrak{g}, \mathfrak{g}))$. Its transpose $r(\text{ad } x)^*$ belongs to $\text{GL}(F(\mathfrak{g}, \mathfrak{g}^*))$ and we have $\beta^*(\omega_{ev, \rho}^i) = r(\text{ad } x)^*(x^i)$. By definition of the Berezinian, the Jacobian J^ρ is equal to $\text{Ber}(r(\text{ad } x)^*)$. The theorem follows from formula (25).

Second proof. We present this less natural proof for fun, and because it is a simplification of the proof of theorem 4.6 below. Let $a \in \mathfrak{g}$. We recall the formula given in [18] for the derivation $\beta^*(-\zeta_a^\lambda)$ of $F(\mathfrak{g})$ obtained by transporting the derivation $-\zeta_a^\lambda$ of $F(G)$. We have

$$(67) \quad \beta^*(-\zeta_a^\lambda) = -\zeta_{p(\text{ad } x)(a)}.$$

Since $p(0) = 1$, it is sufficient to show that for all $a \in \mathfrak{g}$ we have

$$(68) \quad \mathcal{L}(\zeta_{p(\text{ad } x)(a)}) (\text{Ber}(r(\text{ad } x))d) = 0.$$

By (34) and (35) this is equivalent to

$$(69) \quad \zeta_{p(\text{ad } x)(a)} (\text{Ber}(r(\text{ad } x))) + \text{Ber}(r(\text{ad } x)) \text{div}(\zeta_{p(\text{ad } x)(a)}) = 0.$$

By propositions 2.4 and 2.2, this reduces to

$$(70) \quad \text{str}(p(0)w'(\text{ad } x) \text{ad } a) - \text{str}\left(\frac{p(\text{ad } x) - p(0)}{\text{ad } x} \text{ad } a\right) = 0.$$

This identity follows from the relation

$$(71) \quad p(0)w'(t) - \frac{p(t) - p(0)}{t} = 0,$$

which is miraculously true: since $p(0) = 1$, we have

$$p(0)w'(t) = \frac{e^{-t}}{1 - e^{-t}} - \frac{1}{t} = \frac{1}{e^t - 1} - \frac{1}{t} \quad \text{and} \quad \frac{p(t) - p(0)}{t} = \frac{\frac{t}{e^t - 1} - 1}{t} = \frac{1}{e^t - 1} - \frac{1}{t}.$$

□

3. FORMAL HOMOGENEOUS SPACES.

In this section, we consider a Lie superalgebra $\mathfrak{g}$ and a Lie subalgebra $\mathfrak{h} \subset \mathfrak{g}$ which is a direct factor: there exists and we choose a $\mathbb{K}$ -submodule $\mathfrak{q} \subset \mathfrak{g}$ such that

$$(72) \quad \mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}.$$

The following lemma is a straightforward consequence of the fact that β is an isomorphism of coalgebras.

Lemma 3.1. *Let $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{q}$ be a decomposition into a direct sum of submodules. The map $S(\mathfrak{p}) \otimes S(\mathfrak{q}) \ni w \otimes w' \rightarrow \beta(w)\beta(w') \in U(\mathfrak{g})$ is an isomorphism of coalgebras.*

Corollary 3.1. *The map $\bar{\beta} : S(\mathfrak{q}) \otimes U(\mathfrak{h}) \ni w \otimes u \rightarrow \beta(w)u \in U(\mathfrak{g})$ is an isomorphism of coalgebras.*

3.1. Induced and coinduced representations in exponential coordinates. For $a \in \mathfrak{g}$, we recall that the left multiplication l_a is a coderivation of $U(\mathfrak{g})$. We denote by $\bar{l}_a$ the corresponding coderivation of $S(\mathfrak{q}) \otimes U(\mathfrak{h})$. Recall the notation $\Delta_{S(\mathfrak{g})}(w) = \sum w_i \otimes w'_i$ for $w \in S(\mathfrak{q})$.

Proposition 3.1. *Let $a \in \mathfrak{g}$. There exist $\alpha_a \in F(\mathfrak{q}, \mathfrak{q})$ and $\theta_a \in F(\mathfrak{q}, \mathfrak{h})$ such that, for $w \in S(\mathfrak{q})$ and $u \in U(\mathfrak{h})$ we have*

$$(73) \quad \bar{l}_a(w \otimes u) = \sum \pm w_i \alpha_a(w'_i) \otimes u + \sum \pm w_i \otimes j(\theta_a(w'_i))u.$$

Proof. We use the convolution notation for the product in $F(\mathfrak{q}, S(\mathfrak{q}) \otimes U(\mathfrak{h}))$. Define $C \in F(\mathfrak{q}, S(\mathfrak{q}) \otimes U(\mathfrak{h}))$ by

$$(74) \quad C(w) = \bar{l}_a(w \otimes 1) \text{ for } w \in S(\mathfrak{q}).$$

We have $\bar{l}_a(w \otimes u) = C(w)(1 \otimes u)$, and formula (73) is equivalent to

$$(75) \quad C = (\text{id}_{S(\mathfrak{q})} \otimes \text{id}_{U(\mathfrak{h})}) * (\alpha_a \otimes \text{id}_{U(\mathfrak{h})} + \text{id}_{S(\mathfrak{q})} \otimes j \circ \theta_a) = c_{\alpha_a} \otimes 1 + (\text{id}_{S(\mathfrak{q})} \otimes j \circ \theta_a) \circ \Delta_{S(\mathfrak{q})}.$$

To prove (75), we argue as in [20, theorem 1]. We define $\gamma \in F(\mathfrak{q}, S(\mathfrak{q}) \otimes U(\mathfrak{h}))$ by $\gamma = (S \otimes 1) * C$, where $S \in F(\mathfrak{q}, S(\mathfrak{q}))$ is the antipode of $S(\mathfrak{q})$. Since S is the inverse of $\text{id}_{S(\mathfrak{q})}$ for the convolution product, we have $C = (\text{id}_{S(\mathfrak{q})} \otimes 1) * \gamma$. Because $\bar{l}_a$ is a coderivation, γ takes its values in the submodule of primitive elements of $S(\mathfrak{q}) \otimes U(\mathfrak{h})$. Since $\mathbb{K}$ is $\mathbb{Q}$ -algebra, this submodule is $\mathfrak{q} \otimes 1 + 1 \otimes j(\mathfrak{h})$. We write $\gamma = \alpha_a \otimes 1 + 1 \otimes j \circ \theta_a$ with $\alpha_a \in F(\mathfrak{q}, \mathfrak{q})$ and $\theta_a \in F(\mathfrak{q}, \mathfrak{h})$. □

Remark 3.1. Manageable formulas for the maps $\alpha_a \in F(\mathfrak{q}, \mathfrak{q})$ and $\theta_a \in F(\mathfrak{q}, \mathfrak{h})$ are not known in general. In the case $\mathfrak{h} = \{0\}$, a formula for α_a is given in [18] (see (67) above). In the case of symmetric pairs, we give formulas for α_a and θ_a in theorem 4.5.

Let (V, χ) be an $\mathfrak{h}$ -module. We consider the induced representation of $\mathfrak{g}$, that is the representation by left multiplication in $U(\mathfrak{g}) \otimes_{\mathfrak{h}} V$. We define

$$(76) \quad \bar{\beta}_{\chi} : S(\mathfrak{q}) \otimes V \rightarrow U(\mathfrak{g}) \otimes_{\mathfrak{h}} V$$

as in corollary 3.1. For $a \in \mathfrak{g}$, we denote by $l_{\chi,a}$ the endomorphism of $U(\mathfrak{g}) \otimes_{\mathfrak{h}} V$ induced by the l_a , and by $\bar{l}_{\chi,a}$ the corresponding endomorphism of $S(\mathfrak{q}) \otimes V$. It belongs to $F(\mathfrak{q}, S(\mathfrak{q}) \otimes \text{End}(V))$, and it follows from (75) that $\bar{l}_{\chi,a}$ is given by the formula

$$(77) \quad \bar{l}_{\chi,a} : S(\mathfrak{q}) \ni w \mapsto c_{\alpha_a}(w) \otimes \text{id}_V + (\text{id}_{S(\mathfrak{q})} \otimes \chi \circ \theta_a) \circ \Delta_{S(\mathfrak{q})}(w)(1 \otimes \text{id}_V).$$

The coinduced representation is a representation in the space

$$(78) \quad F(G/H, V) := \text{Hom}_{\mathfrak{h}}(U(\mathfrak{g}), V),$$

that is the subspace of elements $g \in F(G, V)$ such that $-\pm g(uj(a)) = a(g(u))$ for $a \in \mathfrak{h}$ and $u \in U(\mathfrak{g})$. The action of $a \in \mathfrak{g}$ in $F(G/H, V)$ is the restriction of $-\zeta_a^{\lambda} \in \text{End}(F(G, V))$, we denote it by $-\zeta_{\chi,a}$.

It follows from corollary 3.1 that the map

$$(79) \quad \bar{\beta}_{\chi}^* : F(G/H, V) \ni \phi \rightarrow \phi \circ \beta \in F(\mathfrak{q}, V)$$

is bijective. We denote by $-\bar{\zeta}_{\chi,a} := -\bar{\beta}_{\chi}^* \circ \zeta_{\chi,a} \circ \bar{\beta}_{\chi}^{*-1}$ the corresponding endomorphism of $F(\mathfrak{q}, V)$. We denote by $\chi(\theta_a)$ the endomorphism $\phi \rightarrow \pm(\chi \circ \theta_a \otimes \phi) \circ \Delta_{S(\mathfrak{q})}$ of $\text{End}_{F(\mathfrak{q})}(F(\mathfrak{q}, V))$. It follows from (77) that

$$(80) \quad -\bar{\zeta}_{\chi,a} = -\zeta_{\alpha_a} + \chi(\theta_a).$$

3.2. The exponential map on a formal homogeneous space. We recall (see formula 78) that $F(G/H)$ is the algebra dual to the coalgebra $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$, i.e. the orthogonal of $U(\mathfrak{g})\mathfrak{h}$ in $F(G)$.

The restriction of δ to $F(G/H)$ is still denoted by δ . It provides $\mathbb{K}$ with a structure of $F(G/H)$ -module. The augmented algebra $F(G/H)$ is thought of as the algebra of functions on the *formal homogeneous space* G/H , and δ is “the evaluation at the point $1 \in G/H$ ”.

Note that $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$ and $F(G/H)$ are the induced and coinduced representations by the trivial representation of $\mathfrak{h}$ in $\mathbb{K}$. We denote this representation by triv , and we use the notations $\tilde{\beta} := \bar{\beta}_{\text{triv}}$ and $\tilde{\beta}^* := \bar{\beta}_{\text{triv}}^*$. In this case,

$$(81) \quad \tilde{\beta} : S(\mathfrak{q}) \rightarrow U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h} \text{ and } \tilde{\beta}^* : F(G/H) \rightarrow F(\mathfrak{q}),$$

are respectively a coalgebra and an algebra isomorphisms. The map $\tilde{\beta}$ is the formal analogue of the exponential map $\mathfrak{q} \rightarrow G/H$.

For $a \in \mathfrak{g}$, we still denote by l_a the left multiplication by $j(a)$ in the quotient $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$, i.e. $l_a = l_{\text{triv},a}$. We denote by ζ_a the transpose map in $F(G/H)$. Thus $\zeta_a = \zeta_{\text{triv},a}$ is the restriction of ζ_a^{λ} to $F(G/H) \subset F(G)$.

Consider the maps in (81). For $a \in \mathfrak{g}$, we denote $\tilde{l}_a := \bar{l}_{\text{triv},a}$ and $-\tilde{\zeta}_a := -\bar{\zeta}_{\text{triv},a}$ the corresponding coderivation of $S(\mathfrak{q})$ and derivation of $F(\mathfrak{q})$. As a particular case of formulas (77) and (80), we have

$$(82) \quad \tilde{l}_a = c_{\alpha_a} \text{ and } -\tilde{\zeta}_a = -\zeta_{\alpha_a}$$

where $\alpha_a \in F(\mathfrak{q}, \mathfrak{q})$ is the vector field defined in proposition 3.1.

3.3. Imprimitivity theorem. The following definition is similar to definition 1.2.

Definition 3.1. A $\mathfrak{g} - F(G/H)$ -module $\mathcal{V}$ is a $\mathbb{K}$ -module provided with a structure of $F(G/H)$ -module and a structure of $\mathfrak{g}$ module which satisfy the Leibnitz relation

$$(83) \quad a(\phi f) = -\zeta_a^\lambda(\phi)f \pm \phi(af), \text{ for } a \in \mathfrak{g}, \phi \in F(G/H), f \in \mathcal{V}.$$

If $\mathcal{V}$ and $\mathcal{W}$ are $\mathfrak{g} - F(G/H)$ -modules, so are $\mathcal{V} \otimes_{F(G/H)} \mathcal{W}$ and $\text{Hom}_{F(G/H)}(\mathcal{V}, \mathcal{W})$.

Induced and coinduced modules by a representation of $\mathfrak{h}$ in a $\mathbb{K}$ -module V are naturally $\mathfrak{g} - F(G/H)$ -modules. Indeed: as $F(G/H, V) \subset F(G, V)$ is stable under multiplication by $F(G/H)$, then $F(G/H, V)$ is a $\mathfrak{g} - F(G/H)$ -module. We still denote by δ the evaluation map $\delta : F(G/H, V) \rightarrow V$. We remark that we have natural isomorphisms of $\mathfrak{g} - F(G/H)$ -modules

$$(84) \quad U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h} \otimes_{F(G/H)} F(G/H, V) \simeq U(\mathfrak{g}) \otimes_{\mathfrak{h}} V,$$

and

$$(85) \quad F(G/H, V) \otimes_{F(G/H)} F(G/H, W) \simeq F(G/H, V \otimes W).$$

The purpose of imprimitivity theorems is to characterize induced and coinduced modules among the $\mathfrak{g} - F(G/H)$ -modules. We state such a theorem for coinduced representations in the particular simple case we need.

Let $\mathcal{V}$ be a $\mathfrak{g} - F(G/H)$ -module. We consider $V_{\mathcal{V}} := \mathcal{V} \otimes_{F(G/H)} \mathbb{K}$ and the corresponding map $\bar{\delta} := \text{id}_{\mathcal{V}} \otimes_{F(G/H)} \delta : \mathcal{V} \rightarrow V_{\mathcal{V}}$. The following straightforward lemma provides $V_{\mathcal{V}}$ with a natural $\mathfrak{h}$ -module structure.

Lemma 3.2. *The kernel of $\bar{\delta}$ in $\mathcal{V}$ is stable under the action of $\mathfrak{h}$.*

Suppose also that V is a free $\mathbb{K}$ -module of finite rank (p, q) . It follows that $F(G/H, V)$ is a free $F(G/H)$ -module of finite rank (p, q) . Moreover, the map $\delta : F(G/H, V) \rightarrow V$ induces an isomorphism of $\mathfrak{h}$ -modules $V_{F(G/H, V)} \rightarrow V$, that is we have $\bar{\delta} = \delta$. In particular,

$$F(G/H, V) \simeq F(G/H, V_{F(G/H, V)}).$$

We need a converse to this property, the imprimitivity theorem, due to Blattner [4] (see also [23]) in a slightly different setting.

Assume that $\mathcal{V}$ is free of finite rank as an $F(G/H)$ -module. The $\mathbb{K}$ -module $V_{\mathcal{V}}$ is free of the same rank. Define a map

$$(86) \quad \tilde{\delta} : \mathcal{V} \rightarrow F(G/H, V_{\mathcal{V}}) \text{ by } \tilde{\delta}(v)(u) = \pm \bar{\delta}(uv).$$

The facts that $\tilde{\delta}$ is well defined, and that it commutes with the actions of $\mathfrak{g}$ and $F(G/H)$ are easy.

Proposition 3.2. *Let $\mathcal{V}$ be a $\mathfrak{g} - F(G/H)$ -module which is free of finite rank as an $F(G/H)$ -module. Then $\tilde{\delta}$ is an isomorphism. In particular*

$$\mathcal{V} \simeq F(G/H, V_{\mathcal{V}}).$$

Proof. Let (ϕ_i) be a basis of $\mathcal{V}$, and $v_i := \bar{\delta}(\phi_i) \in V_{\mathcal{V}}$. Then (v_i) is a basis of $V_{\mathcal{V}}$. We have $\delta(\tilde{\delta}(\phi_i)) = v_i$. It follows from lemma 1.2 that $(\tilde{\delta}(\phi_i))$ is a basis of $F(G/H, V_{\mathcal{V}})$. $\square$

3.4. Dualizing modules on formal homogeneous spaces. In this section, we assume that the $\mathbb{K}$ -module $\mathfrak{q} = \mathfrak{g}/\mathfrak{h}$ is free of finite rank (p, q) .

By (81), the algebra $F(G/H)$ is formally smooth of dimension (p, q) and the dualizing module $K(F(G/H))$ is defined. By definition, it is free of rank 1 as an $F(G/H)$ -module. For $a \in \mathfrak{g}$, $-\zeta_a^\lambda$ is a derivation of $F(G/H)$. It acts in $K(F(G/H))$, so $K(F(G/H))$ is in a natural way a $\mathfrak{g} - F(G/H)$ -module (recall definition 3.1). The evaluation

$$(87) \quad \bar{\delta} : K(F(G/H)) \rightarrow K(F(G/H)) \otimes_{F(G/H)} \mathbb{K}$$

is a morphism of $\mathfrak{h}$ -modules. It follows that the $\mathbb{K}$ -isomorphism $K(F(G/H)) \otimes_{F(G/H)} \mathbb{K} \simeq \text{Ber}((\mathfrak{g}/\mathfrak{h})^*)$ given in (32) is an isomorphism of $\mathfrak{h}$ -modules. It follows from proposition 3.2 that the map $\tilde{\delta}$ defined in (86) is an isomorphism

$$(88) \quad \tilde{\delta} : K(F(G/H)) \rightarrow F(G/H, \text{Ber}((\mathfrak{g}/\mathfrak{h})^*)).$$

of $\mathfrak{g} - F(G/H)$ -modules.

For $a \in \mathfrak{h}$, $\text{ad } a$ induces an endomorphism of $\mathfrak{g}/\mathfrak{h}$. We denote by $\text{str}_{\mathfrak{g}/\mathfrak{h}}(\text{ad } a)$ its supertrace. The action of a in $\text{Ber } \mathfrak{g}/\mathfrak{h}$ is the multiplication by $\text{str}_{\mathfrak{g}/\mathfrak{h}}(\text{ad } a)$, and the action of a in $\text{Ber}((\mathfrak{g}/\mathfrak{h})^*)$ is the multiplication by $-\text{str}_{\mathfrak{g}/\mathfrak{h}}(\text{ad } a)$.

Proposition 3.3. *The dualizing module $K(F(G/H))$ has a $\mathfrak{g}$ -invariant basis as an $F(G/H)$ -module if and only if the unimodularity condition (10) holds. In this case, the $\mathbb{K}$ -submodule of $\mathfrak{g}$ -invariant elements of $K(F(G/H))$ is free of rank one, and a basis of the $\mathbb{K}$ -module of $\mathfrak{g}$ -invariant is a basis of the $F(G/H)$ -module $K(F(G/H))$.*

Proof. Let D be an invariant basis of $K(F(G/H))$. The element $\tilde{\delta}(D)$ is a basis of $\text{Ber}(\mathfrak{g}/\mathfrak{h})^*$ invariant by $\mathfrak{h}$. So (10) holds.

Conversely, suppose that (10) holds. Then $K(F(G/H))$ is isomorphic to $F(G/H)$ —see (88)—, and the constant function $1 \in F(G/H)$ is an invariant basis of $F(G/H)$. $\square$

Remark 3.2. *Similarly, the $\mathfrak{g} - F(G/H)$ -module $\mathcal{T}(F(G/H))$ of derivations of $F(G/H)$ is isomorphic to $F(G/H, \mathfrak{g}/\mathfrak{h})$ and the module $\Omega_{ev}^1(F(G/H))$ to $F(G/H, (\mathfrak{g}/\mathfrak{h})^*)$. However, even under assumption (10), there is usually no basis of $F(G/H, \mathfrak{g}/\mathfrak{h})$ consisting of $\mathfrak{g}$ -invariant elements. So the simple proof of proposition 3.3, given in proposition 2.6 for the special case $\mathfrak{h} = 0$, does not generalize.*

To get invariant elements without unimodularity hypotheses, we consider twisted dualizing modules. Let V be an $\mathfrak{h}$ -module. We define

$$(89) \quad K(F(G/H), V) := K(F(G/H)) \otimes_{F(G/H)} F(G/H, V),$$

and we call it the *dualizing module twisted by V* . As in (88), there is a canonical isomorphism of (recall (85) and (88)) $\mathfrak{g} - F(G/H)$ -modules

$$(90) \quad K(F(G/H), V) \simeq F(G/H, \text{Ber}((\mathfrak{g}/\mathfrak{h})^*) \otimes_{\mathbb{K}} V).$$

We apply this construction to the $\mathfrak{h}$ -module $\text{Ber } \mathfrak{g}/\mathfrak{h}$. Using the isomorphism (26), we obtain a canonical isomorphism of $\mathfrak{g} - F(G/H)$ -modules

$$(91) \quad K(F(G/H), \text{Ber } \mathfrak{g}/\mathfrak{h}) \simeq F(G/H).$$

We denote by $\mathbf{D}_{G/H}$ the element of $K(F(G/H), \text{Ber } \mathfrak{g}/\mathfrak{h})$ which corresponds to $1 \in F(G/H)$ by this isomorphism.

Proposition 3.4. *i) The element $\mathbf{D}_{G/H} \in K(F(G/H), \text{Ber } \mathfrak{g}/\mathfrak{h})$ is $\mathfrak{g}$ -invariant and $\tilde{\delta}(\mathbf{D}_{G/H}) \in \text{Ber}(\mathfrak{g}/\mathfrak{h})^* \otimes_{\mathbb{K}} \text{Ber } \mathfrak{g}/\mathfrak{h}$ is equal to the canonical element $\mathbf{d}$ defined in (27).*

ii) The $\mathbb{K}$ -module of $\mathfrak{g}$ -invariant elements of $K(F(G/H), \text{Ber } \mathfrak{g}/\mathfrak{h})$ is free of rank one with basis $\mathbf{D}_{G/H}$.

iii) $\mathbf{D}_{G/H}$ is a basis of the $F(G/H)$ -module $K(F(G/H), \text{Ber } \mathfrak{g}/\mathfrak{h})$.

3.5. The Jacobian of the exponential map on a formal homogeneous space. We keep the assumptions of subsection 3.4. The isomorphism $\tilde{\beta}^* : F(G/H) \rightarrow F(\mathfrak{q})$ of augmented algebras—see (81)—induces like in (37) a $\mathbb{K}$ -isomorphism $\tilde{\beta}^* : K(F(G/H)) \rightarrow K(F(\mathfrak{q}))$. Since the natural action of $a \in \mathfrak{g}$ in $F(G/H)$ is given by the derivation $-\zeta_a$ (see subsection 3.2), the natural action in $K(G/H)$ is given by $-\mathcal{L}(\zeta_a)$. By (82) and (39), the natural action of a in $K(F(\mathfrak{q}))$ obtained by transporting $-\mathcal{L}(\zeta_a)$ by $\tilde{\beta}^*$ is equal to $-\mathcal{L}(\zeta_{\alpha_a})$.

We recall that χ is a representation of $\mathfrak{h}$ in V . We define

$$(92) \quad K(F(\mathfrak{q}), V) := K(F(\mathfrak{q})) \otimes_{F(\mathfrak{q})} F(\mathfrak{q}, V).$$

Because of (38), the natural map $\tilde{\beta}^* \otimes \bar{\beta}_\chi^* : K(F(G/H)) \otimes_{\mathbb{K}} F(G/H, V) \rightarrow K(F(\mathfrak{q}), V)$ induces an isomorphism $\bar{\beta}_\chi^* : K(F(G/H), V) \rightarrow K(F(\mathfrak{q}), V)$. Recall that the action of a in $F(\mathfrak{q}, V)$ is denoted by $-\tilde{\zeta}_{\chi, a}$. It follows that the action of a in $K(F(\mathfrak{q}), V)$ is given by $-\mathcal{L}(\zeta_{\alpha_a}) \otimes 1 - 1 \otimes \tilde{\zeta}_{\chi, a}$. From (80) we get the following formula for the action of $a \in \mathfrak{g}$ in $K(F(\mathfrak{q}), V)$:

$$(93) \quad a(d \otimes \phi) = -\mathcal{L}(\zeta_{\alpha_a})(d) \otimes \phi \pm (-d \otimes \zeta_{\alpha_a}(\phi) + d \otimes \chi(\theta_a)(\phi)) \text{ for } d \in K(F(\mathfrak{q})), \phi \in F(\mathfrak{q}, V).$$

We choose a basis $d_{\mathfrak{q}}$ of $K(F(\mathfrak{q}))$ which is invariant by translations (see lemma 1.5). An element of $K(F(\mathfrak{q}), V)$ can be written in an unique manner as $d_{\mathfrak{q}}\phi$, with $\phi \in F(\mathfrak{q}, V)$ —we leave out the symbol $\otimes$. Formula (93) specializes (see (35) and (36)) to

$$(94) \quad a(d_{\mathfrak{q}}\phi) = \pm d_{\mathfrak{q}}(-\operatorname{div}(\zeta_{\alpha_a})\phi - \zeta_{\alpha_a}(\phi) + \chi(\theta_a)(\phi)) \text{ for } a \in \mathfrak{g}, \phi \in F(\mathfrak{q}, V).$$

We apply these considerations to the $\mathfrak{h}$ -module $V = \operatorname{Ber} \mathfrak{g}/\mathfrak{h}$. We recall (see subsection 3.4) that the action of $\mathfrak{h}$ is the multiplication by the linear form $\chi = \operatorname{str}_{\mathfrak{g}/\mathfrak{h}}$. We denote by

$$(95) \quad \mathbf{d}_{\mathfrak{q}} \in K(F(\mathfrak{q}), \operatorname{Ber} \mathfrak{g}/\mathfrak{h})$$

the canonical element, that is, recalling the definition (27) the element such that $\delta(\mathbf{d}_{\mathfrak{q}}) = \mathbf{d} \in \operatorname{Ber}(\mathfrak{g}/\mathfrak{h})^* \otimes \operatorname{Ber} \mathfrak{g}/\mathfrak{h}$.

An element of $K(F(\mathfrak{q}), \operatorname{Ber} \mathfrak{g}/\mathfrak{h})$ can be written in a unique manner as $\phi \mathbf{d}_{\mathfrak{q}}$, with $\phi \in F(\mathfrak{q})$. From equation (94) we get

$$(96) \quad a(\phi \mathbf{d}_{\mathfrak{q}}) = (-\operatorname{div}(\zeta_{\alpha_a})\phi - \zeta_{\alpha_a}(\phi) + \operatorname{str}_{\mathfrak{g}/\mathfrak{h}} \circ \theta_a \phi) \mathbf{d}_{\mathfrak{q}} \text{ for } a \in \mathfrak{g}, \phi \in F(\mathfrak{q}).$$

We defined $\mathbf{D}_{G/H} \in K(F(G/H), \operatorname{Ber} \mathfrak{g}/\mathfrak{h})$ in proposition 3.4. There exists a function $J \in F(\mathfrak{q})$ such that

$$(97) \quad \bar{\beta}_{\operatorname{str}_{\mathfrak{g}/\mathfrak{h}}}^*(\mathbf{D}_{G/H}) = J \mathbf{d}_{\mathfrak{q}}.$$

We call J the *Jacobian of the exponential map* $\mathfrak{q} \rightarrow G/H$. A natural problem is to compute J . We already did it in theorem 2.1 when $\mathfrak{h} = 0$. In section 4.3 we compute J in the case of symmetric pairs. We shall use the following characterization of J , which follows immediately from (96).

Lemma 3.3. *The function $J \in F(\mathfrak{q})$ is the unique element of $F(\mathfrak{q})_{\mathbf{0}}$ which satisfies $\delta(J) = 1$ and such that, for all $a \in \mathfrak{g}$,*

$$(98) \quad \frac{\zeta_{\alpha_a}(J)}{J} + \operatorname{div}(\zeta_{\alpha_a}) - \operatorname{str}_{\mathfrak{g}/\mathfrak{h}} \circ \theta_a = 0.$$

Remark 3.3. *By proposition 3.3, if the unimodularity condition (10) holds, there exists a $\mathfrak{g}$ -invariant basis $D_{G/H}$ of $K(F(G/H))$. If we ask also for the equality $\bar{\delta}(D_{G/H}) = \bar{\delta}(d_{\mathfrak{q}})$ in $\operatorname{Ber}(\mathfrak{q}^*)$, it is unique, and we obtain, with the same function J than in (97),*

$$(99) \quad \tilde{\beta}^*(D_{G/H}) = J d_{\mathfrak{q}}.$$

3.6. Homogeneous spaces of purely odd dimension. We keep the notations of subsection 3.4, and in addition we assume that $\dim \mathfrak{g}/\mathfrak{h} = (0, q)$. We apply the results of subsection 1.6 to the algebra $F(G/H)$.

Since it is free of finite rank, we have $F(G/H)_* = F(G/H)^* = U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$, and Berezin integral provides a canonical isomorphism $\psi_{F(G/H)} : K(F(G/H)) \rightarrow U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$ of $\mathfrak{g} - F(G/H)$ -modules (see subsection 1.6). Let V be an $\mathfrak{h}$ -module. Tensoring with $F(G/H, V)$, we obtain a canonical isomorphism $K(F(G/H), V) \rightarrow U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h} \otimes_{F(G/H)} F(G/H, V)$. Composing with the isomorphism (84), we obtain a canonical isomorphism of $\mathfrak{g} - F(G/H)$ -modules

$$(100) \quad \kappa_\chi : K(F(G/H), V) \simeq U(\mathfrak{g}) \otimes_{\mathfrak{h}} V,$$

and, using (90), a canonical isomorphism of $\mathfrak{g} - F(G/H)$ -modules

$$(101) \quad F(G/H, \text{Ber}(\mathfrak{g}/\mathfrak{h})^* \otimes V) \simeq U(\mathfrak{g}) \otimes_{\mathfrak{h}} V.$$

Remark 3.4. *The isomorphism (101) between an induced and a coinduced representation of $\mathfrak{g}$ is well known, see [2, 5–7]. It is used in Gorelik’s paper [9]. The properties we need in this article, that is the relation with dualizing modules and the fact that it is an isomorphism of $F(G/H)$ -modules, are explicit in [5–7].*

In particular, using the canonical isomorphism (26), the $\mathfrak{g} - F(G/H)$ -module $U(\mathfrak{g}) \otimes_{\mathfrak{h}} \text{Ber } \mathfrak{g}/\mathfrak{h}$ is canonically isomorphic to $F(G/H)$. We denote by $\mathbf{T} \in U(\mathfrak{g}) \otimes_{\mathfrak{h}} \text{Ber } \mathfrak{g}/\mathfrak{h}$ the element corresponding to $1 \in F(G/H)$. Remember (see proposition 3.4) the definition of the canonical element $\mathbf{D}_{G/H} \in K(F(G/H), \text{Ber } \mathfrak{g}/\mathfrak{h})$. We have

$$(102) \quad \kappa_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}(\mathbf{D}_{G/H}) = \mathbf{T}.$$

We have proved the following proposition, analogous to proposition 3.4.

Proposition 3.5. *a) The $\mathbb{K}$ -module of $\mathfrak{g}$ -invariants of $U(\mathfrak{g}) \otimes_{\mathfrak{h}} \text{Ber } \mathfrak{g}/\mathfrak{h}$ is free of rank 1 with basis $\mathbf{T}$. Moreover, $\mathbf{T}$ is a basis of the $F(G/H)$ -module $U(\mathfrak{g}) \otimes_{\mathfrak{h}} \text{Ber } \mathfrak{g}/\mathfrak{h}$, and the map $F(G/H) \ni \phi \rightarrow \mathbf{T}\phi \in U(\mathfrak{g}) \otimes_{\mathfrak{h}} \text{Ber } \mathfrak{g}/\mathfrak{h}$ is an isomorphism of $\mathfrak{g} - F(G/H)$ -modules.*

b) Assume condition (10). Then the $\mathbb{K}$ -module of $\mathfrak{g}$ -invariants of $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$ is free of rank 1. A basis T of this module is also a basis of the $F(G/H)$ -module $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$. The map $\phi \rightarrow T\phi$ from $F(G/H)$ to $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$ is an isomorphism of $\mathfrak{g} - F(G/H)$ -modules.

Similarly, we have the isomorphism of $\mathcal{T}(F(\mathfrak{q})) - F(\mathfrak{q})$ -modules $\psi_{F(\mathfrak{q})} : K(F(\mathfrak{q})) \rightarrow S(\mathfrak{q})$, and lemma 1.7 gives

$$(103) \quad \psi_{F(G/H)} = \tilde{\beta} \circ \psi_{F(\mathfrak{q})} \circ \tilde{\beta}^*.$$

Tensoring with $F(\mathfrak{q}, V)$, we obtain a canonical isomorphism

$$(104) \quad \bar{\kappa}_{\chi} : K(F(\mathfrak{q}), V) \rightarrow S(\mathfrak{q}) \otimes_{F(\mathfrak{q})} F(\mathfrak{q}, V) \simeq S(\mathfrak{q}) \otimes V.$$

Formula (103) extends to

$$(105) \quad \kappa_{\chi} = \bar{\beta}_{\chi} \circ \bar{\kappa}_{\chi} \circ \bar{\beta}_{\chi}^*.$$

We identify $\mathfrak{q}^*$ and $(\mathfrak{g}/\mathfrak{h})^*$. Since the rank of $\mathfrak{g}/\mathfrak{h}$ is $(0, q)$, we have $\text{Ber } \mathfrak{g}/\mathfrak{h} = S^q(\mathfrak{q}^*)$. There is a canonical element $\mathbf{t} \in S(\mathfrak{q}) \otimes \text{Ber } \mathfrak{g}/\mathfrak{h}$: in term of a totally ordered basis $(e_i)_{1 \leq i \leq q}$ of $\mathfrak{q}$ and of its dual basis $(x^i)_{1 \leq i \leq q}$ of $\mathfrak{q}^*$, we have

$$(106) \quad \mathbf{t} = (-1)^q e_1 \cdots e_q \otimes x^q \cdots x^1.$$

Recall the definition (95) of $\mathbf{d}_{\mathfrak{q}} \in K(F(\mathfrak{q}), \text{Ber } \mathfrak{g}/\mathfrak{h})$. The analogue of (102) is the formula

$$(107) \quad \bar{\kappa}_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}(\mathbf{d}_{\mathfrak{q}}) = \mathbf{t}.$$

Recall the Jacobian $J \in F(\mathfrak{q})$ of the exponential map $\mathfrak{q} \rightarrow G/H$ (formula (97)). *The following theorem explains the role of this Jacobian in the study of Gorelik’s elements.*

Theorem 3.1. *a) We have $\bar{\beta}_{\text{str}}(J\mathbf{t}) = \mathbf{T}$.*

b) Suppose that condition (10) holds. Then $\tilde{\beta}(J e_1 \cdots e_q) \in U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$ is a basis of the $\mathbb{K}$ -module of $\mathfrak{g}$ -invariants of $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$, and a basis of $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$ as a $F(G/H)$ -module.

Proof. a) From (102) and (97) we get

$$\bar{\beta}_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}^*(\kappa_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}^{-1}(\mathbf{T})) = J\mathbf{d}_{\mathfrak{q}}.$$

Using the fact that $\bar{\kappa}_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}$ is a $F(\mathfrak{q})$ -module isomorphism, we obtain

$$\bar{\kappa}_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}(\bar{\beta}_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}^*(\kappa_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}^{-1}(\mathbf{T}))) = J\bar{\kappa}_{\text{str}_{\mathfrak{g}/\mathfrak{h}}}(\mathbf{d}_{\mathfrak{q}}).$$

We get our formula using (107) and (105).

b) Follows from part a) and proposition 3.5. $\square$

4. SYMMETRIC PAIRS.

In this section, we consider a Lie symmetric pair $(\mathfrak{g}, \sigma)$ with corresponding decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$.

4.1. Representations by coderivations. Let $a \in \mathfrak{g}$. Recall the coderivation $\tilde{l}_a = c_{\alpha_a}$ of $S(\mathfrak{q})$ (formula (82)). In this subsection, we give a formula for the vector field $\alpha_a \in F(\mathfrak{q}, \mathfrak{q})$. In fact, we do more. Following the method of [18], we describe *all the representations by coderivations of $\mathfrak{g}$ in $S(\mathfrak{q})$ which are universal*. Informally, this means that they are given by formulas which are the same for all symmetric pairs $(\mathfrak{g}, \sigma)$. Formally, it can be express in functorial terms (see [18]). Reasoning as in [18], these representations are of a very special form, and we are led to the following considerations.

We denote by $y := x_{\mathfrak{q}} \in F(\mathfrak{q}, \mathfrak{q})$ the generic point of $\mathfrak{q}$. We consider y as the element of the Lie $F(\mathfrak{q})$ -superalgebra $F(\mathfrak{q}, \mathfrak{g})$ which is 0 on $S^n(\mathfrak{q})$ for $n \neq 1$, and which is the natural injection $\mathfrak{q} \rightarrow \mathfrak{g}$ on $\mathfrak{q}$. The decomposition $F(\mathfrak{q}, \mathfrak{g}) = F(\mathfrak{q}, \mathfrak{h}) \oplus F(\mathfrak{q}, \mathfrak{q})$ is a symmetric pair in the category of Lie $F(\mathfrak{q})$ -superalgebras. For a formal power series $p(t) \in \mathbb{K}_0[[t]]$ the element $p(\text{ad } y) \in \text{End}_{F(\mathfrak{q})}(F(\mathfrak{q}, \mathfrak{g}))$ is defined by locally finite sums like in (17). In particular, for $a \in \mathfrak{g}$, the element $p(\text{ad } y)(a) \in F(\mathfrak{q}, \mathfrak{g})$ is defined by the following formula, similar to (49). Let $b_1, b_2, \dots, b_n$ in $\mathfrak{q}$. We have

$$(108) \quad p(\text{ad } y)(a) : b_1 \cdots b_n \mapsto p_n \sum_{s \in S_n} \pm \text{ad } b_{s(1)} \cdots \text{ad } b_{s(n)}(a).$$

We consider two series $d(t), p(t) \in \mathbb{K}_0[[t]]$ such that $p(t) = p(-t)$ and $d(t) = -d(-t)$. We write $p(t) = p_0 + p_2 t^2 + \cdots \in \mathbb{K}_0[[t^2]]$ and $d(t) = d_1 t + d_3 t^3 + \cdots \in t\mathbb{K}_0[[t^2]]$. Formula (108) implies that for $a \in \mathfrak{q}$ we have $p(\text{ad } y)(a) \in F(\mathfrak{q}, \mathfrak{q})$, and for $a \in \mathfrak{h}$ we have $d(\text{ad } y)(a) \in F(\mathfrak{q}, \mathfrak{q})$. In particular, the following definition is meaningful.

Definition 4.1. For $a \in \mathfrak{q}$, we denote by C^a (or $C_{p,d,\mathfrak{g},\sigma}^a$ if we need to specify) the coderivation $c_{p(\text{ad } y)(a)}$ of $F(\mathfrak{q})$. For $a \in \mathfrak{h}$, we denote by C^a the coderivation $c_{d(\text{ad } y)(a)}$ of $F(\mathfrak{q})$. For $a \in \mathfrak{g}$, we define C^a by additivity.

The bracket $[C^a, C^b]$ has been computed in [18]. Reasoning as in [18], we obtain the following theorem.

Theorem 4.1. The map $a \rightarrow C_{p,d,\mathfrak{g},\sigma}^a$ is a representation of $\mathfrak{g}$ in $S(\mathfrak{q})$ for any symmetric pair $(\mathfrak{g}, \sigma)$ if and only if the following functional equations hold in $\mathbb{K}[[t, u]]$:

$$(109) \quad d(u) \frac{d(t+u) - d(t)}{u} + d(t) \frac{d(t+u) - d(u)}{t} = -d(t+u),$$

$$(110) \quad p(u) \frac{p(t+u) - p(t)}{u} + p(t) \frac{p(t+u) - p(u)}{t} = -d(t+u),$$

$$(111) \quad p(u) \frac{d(t+u) - d(t)}{u} + d(t) \frac{p(t+u) - p(u)}{t} = -p(t+u).$$

Even if we know it in advance, it is remarkable that these functional equations have non trivial solutions.

Lemma 4.1. Let (p, d) be a solution of the equations of theorem 4.1.

If $d = 0$ or $p = 0$, then $(p, d) = (0, 0)$.

Assume $d \neq 0$. If the first non zero coefficient of d is not a divisor of 0 in $\mathbb{K}_0$, then $d = -t$ and $p_0 \neq 0$.

Proof. The first assertion follows from formulas (111) and (110).

Formula (109) evaluated at $u = 0$ gives $d(t) \left(\frac{d(t)}{t} + 1 \right) = 0$. If the first non zero coefficient of d is not a divisor of 0 in $\mathbb{K}_0$, the multiplication by d is injective, and we get $\frac{d(t)}{t} + 1 = 0$.

We now assume that $d = -t$. Equations (109) and (111) are satisfied, and we are left with the functional equation

$$(112) \quad p(u) \frac{p(t+u) - p(t)}{u} + p(t) \frac{p(t+u) - p(u)}{t} = t + u.$$

Evaluating at $u = 0$ we obtain

$$(113) \quad p_0 p'(t) + p(t) \frac{p(t) - p_0}{t} = t.$$

Suppose $p_0 = 0$. This gives $p(t)^2 = t^2$. Since $p(t) \in t^2 \mathbb{K}[[t^2]]$, this is impossible. $\square$

Lemma 4.1 implies that all the non zero solutions of the functional equations of theorem 4.1 with coefficient in a field $k \subset \mathbb{K}_0$ are given by theorem 4.2 below. We recall that the Bernoulli numbers $b_n \in \mathbb{Q}$ are defined by the generating series $\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} b_n \frac{t^n}{n!}$. For $c \in \mathbb{K}_0^\times$, we define $p_c(t) \in \mathbb{K}_0[[t^2]]$ by

$$(114) \quad p_c(t) = t \coth\left(\frac{t}{c}\right) = c + \sum_{n=1}^{\infty} b_{2n} \frac{2^{2n} t^{2n}}{c^{2n-1} (2n)!} = c + \frac{1}{3c} t^2 - \frac{1}{45c^3} t^4 + \dots.$$

Theorem 4.2. *Let $c \in \mathbb{K}_0^\times$. There exists an unique element $p(t) \in \mathbb{K}_0[[t^2]]$ such that $p(0) = c$ and such that the pair $(p(t), -t)$ is a solution of the functional equations of theorem 4.1. We have $p(t) = p_c(t)$.*

Proof. We repeat the arguments of [18]. The pair $(p(t), -t)$ is a solution if and only if the pair $(p(\frac{t}{c})c, -t)$ is a solution. So we assume that $p_0 = 1$. In this case, according to formula (113), p will be a solution of the (singular) Cauchy problem

$$(115) \quad t p'(t) + p(t)(p(t) - 1) = t^2 \text{ and } p(0) = 1.$$

Writing (we forget for a moment that p is supposed to be even) $p(t) = 1 + p_1 t + p_2 t^2 + \dots$, we get inductively $p_1 = 0$, $p_2 = \frac{1}{3}$, and p_n in function of the p_k with $k < n$. This proves that there is at most one solution of (115).

We leave to the reader to check that the function $t \coth(t)$ is indeed a solution of the differential equation (115), and — this is much stronger — of the functional equation (112). $\square$

Remark 4.1. *As in [18], theorem 4.1 is meaningful and true without assuming that $\mathbb{K}_0$ contains $\mathbb{Q}$, but only assuming that 2 is invertible in $\mathbb{K}$. However, the functional equations have no solutions if $\mathbb{K}_0$ does not contains $\mathbb{Q}$.*

Let $c \in \mathbb{K}_0^\times$. We denote by $a \rightarrow \alpha_c^a$ the linear map from $\mathfrak{g}$ to the space of $F(\mathfrak{q}, \mathfrak{q})$ of vector fields on $\mathfrak{q}$ such that

$$(116) \quad \alpha_c^a = [a, y] \text{ for } a \in \mathfrak{h}, \text{ and } \alpha_c^a = p_c(\text{ad } y)(a) \text{ for } a \in \mathfrak{q}.$$

We denote by

$$(117) \quad C_c^a := c_{\alpha_c^a}$$

the corresponding coderivation of $S(\mathfrak{q})$. It follows from theorem 4.1 and theorem 4.2 that $C_c : a \rightarrow C_c^a$ is a representation of $\mathfrak{g}$ in $S(\mathfrak{q})$.

Remark 4.2. *We denote by I_c the automorphism of the Hopf algebra $S(\mathfrak{q})$ which is the multiplication by c^n in $S^n(\mathfrak{q})$. It intertwines the representations C_1 and C_c in $S(\mathfrak{q})$. Thus, if $\mathbb{K}$ is a field of characteristic 0, there are, up to equivalence, exactly two universal representations by coderivations of $\mathfrak{g}$ in $S(\mathfrak{q})$: the trivial one and C_1 .*

Remark 4.3. *In term of representations, the equality $d = -t$ means the following. Let $a \in \mathfrak{h}$. Then C_c^a (which does not depend on c) is a derivation of $S(\mathfrak{q})$ (of course, by definition, it is also a coderivation). It is determined by the fact that*

$$(118) \quad C_c^a(b) = [a, b] \text{ for } b \in \mathfrak{q}.$$

Thus, C_c^a is the natural adjoint action in of $a \in \mathfrak{h}$ in $S(\mathfrak{q})$.

For $a \in \mathfrak{q}$ and $w \in S^n(\mathfrak{q})$, formula (114) gives an explicit expression for $C_c^a(w) \in S(\mathfrak{q})$ in terms of the Bernoulli numbers and of the coproduct of w . We give a few terms. For $b, b_1, b_2 \in \mathfrak{q}$, we have

$$(119) \quad C_c^a(1) = ca, \quad C_c^a(b) = cab, \quad C_c^a(b_1b_2) = cab_1b_2 + \frac{1}{3c}([a, b_1], b_2] \pm [[a, b_2], b_1]);$$

and for $n > 2$ and $b_1, \dots, b_n \in \mathfrak{q}$, we have

$$(120) \quad C_c^a(b_1 \cdots b_n) = cab_1 \cdots b_n + \frac{1}{3c} \sum_{1 \leq i < j \leq n} \left(\pm [[a, b_i], b_j] \pm [[a, b_j], b_i] \right) b_1 \cdots \hat{b}_i \cdots \hat{b}_j \cdots b_n + \dots$$

where the hat symbol means omission, and the non written terms are of degree $< n - 2$.

4.1.1. *Formulas for and $\tilde{l}_a$ and α_a .* The map $a \rightarrow C_c^a$ extends to an homomorphism of algebras $u \rightarrow C_c^u$ from $U(\mathfrak{g})$ to $\text{End}(S(\mathfrak{q}))$. For $u \in U(\mathfrak{g})$ we write $\tilde{\tau}(u) = C_1^u(1)$. This defines a map $\tilde{\tau} : U(\mathfrak{g}) \rightarrow S(\mathfrak{q})$ which intertwines the left action of $\mathfrak{g}$ in $U(\mathfrak{g})$ and the representation C_1 . Since $C_1^a(1) = 0$ for $a \in \mathfrak{h}$, this gives an intertwining map

$$(121) \quad \tau : U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h} \rightarrow S(\mathfrak{q}).$$

The map τ can be explicitly written in terms of Bernoulli numbers. For example, for b, b_1, b_2, b_3 in $\mathfrak{q}$ we have

$$(122) \quad \tau(1) = 1, \quad \tau(j(b)) = b, \quad \tau(j(b_1)j(b_2)) = b_1b_2,$$

$$(123) \quad \tau(j(b_1)j(b_2)j(b_3)) = b_1b_2b_3 + \frac{1}{3}([b_1, b_2], b_3] \pm [[b_1, b_3], b_2]).$$

For the following theorem, recall the notations α_a and $\tilde{l}_a$ of proposition 3.1 and the notation α_1^a of (116).

Theorem 4.3. *i) The map τ is the inverse of the map $\tilde{\beta} : S(\mathfrak{q}) \rightarrow U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$.*

ii) For $a \in \mathfrak{g}$, we have $\tilde{l}_a = C_1^a$

iii) For $a \in \mathfrak{g}$, we have $\alpha_a = \alpha_1^a$.

Proof. The proof of part i) is similar to the proof of Poincaré-Birkhoff-Witt theorem in [18]. Here is a brief sketch. Let $b \in \mathfrak{q}_0$ and $n \in \mathbb{N}$. By induction on n , we obtain that $\tau(j(b)^n) = b^n$. Applying this to the generic point y of $\mathfrak{q}$ in the Lie superalgebra $F(\mathfrak{q}, \mathfrak{g})$ proves part a). Part ii) follows from part i), and part iii) is a rephrasing of ii). $\square$

Remark 4.4. *When $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is a Lie superalgebra over a field of characteristic zero, and $\mathfrak{h} = \mathfrak{g}_0$, the formula $\tilde{l}_a = C_1^a$ is due to Koszul [13]. Presumably, his proof extends to the general case treated in theorem 4.3. However, we think that our method of proof is interesting: it says that the properties of Bernoulli numbers which make the theorem true are encoded by the fact that $t \coth(t)$ satisfies the functional equation (112).*

4.2. Induced and coinduced representations. We keep the notations of subsection 4.1. We consider in addition a representation $\chi : \mathfrak{h} \rightarrow \text{End}(V)$ of $\mathfrak{h}$ in a $\mathbb{K}$ -module V .

Let $a \in \mathfrak{g}$. We give an explicit formula for $\bar{l}_{\chi,a}$ defined in (77). We follow a method similar to the one of subsection 4.1, where we consider the case of the trivial representation of $\mathfrak{h}$ in $\mathbb{K}$.

Let (h, q) be a pair of formal series $h = h_0 + h_2 t^2 + \dots \in \mathbb{K}_0[[t^2]]$ and $q(t) = q_1 t + q_3 t^3 + \dots \in t\mathbb{K}_0[[t^2]]$. We define a linear map $\mathfrak{g} \ni a \rightarrow \theta^a \in F(\mathfrak{q}, \mathfrak{h})$ by the formulas

$$(124) \quad \theta^a = h(\text{ad } y)(a) \text{ for } a \in \mathfrak{h}, \text{ and } \theta^a = q(\text{ad } y)(a) \text{ for } a \in \mathfrak{q}.$$

Let $c \in \mathbb{K}_0^\times$, and recall the representation C_c of $\mathfrak{g}$ in $S(\mathfrak{q})$ (see (117)). Following the model (77), we define an endomorphism $\Theta_\chi^a \in \text{End}(S(\mathfrak{q}) \otimes V)$ (or $\Theta_{c,\chi,h,q,\mathfrak{g},\sigma}^a$ if we need to specify) by the following formula:

$$(125) \quad \Theta_\chi^a(w \otimes v) = C_c^a(w) \otimes v + (\text{id}_{S(\mathfrak{q})} \otimes \chi \circ \theta^a) \circ \Delta_{S(\mathfrak{q})}(w)(1 \otimes v).$$

For example, for $a \in \mathfrak{h}$, we have:

$$(126) \quad \Theta_\chi^a(1 \otimes v) = 1 \otimes \chi(h_0 a)(v).$$

We denote by Θ_χ the map $a \rightarrow \Theta_\chi^a$, and we study when it is a representation.

Lemma 4.2. *i) The map Θ_χ is a representation of $\mathfrak{g}$ if and only if*

$$\chi \circ (\theta^a \circ C_c^b - \pm \theta^b \circ C_c^a + [\theta^a, \theta^b] - \theta^{[a,b]}) = 0 \quad \forall a, b \in \mathfrak{g}.$$

ii) The map Θ_χ is a representation of $\mathfrak{g}$ for all representations χ of $\mathfrak{h}$ if and only if

$$\theta^a \circ C_c^b - \pm \theta^b \circ C_c^a + [\theta^a, \theta^b] - \theta^{[a,b]} = 0 \quad \forall a, b \in \mathfrak{g}.$$

Proof. We write $\Theta = \Theta_\chi$, $C = C_c$, and $\Delta = \Delta_{S(\mathfrak{q})}$ for simplicity.

i) Let $w \otimes v \in S(\mathfrak{q}) \otimes V$ and $a, b \in \mathfrak{g}$. As C^a is a coderivation we get

$$\begin{aligned} & \Theta^a \circ \Theta^b(w \otimes v) = \\ &= C^a \circ C^b(w) \otimes v + (C^a \otimes \chi \circ \theta^b \pm C^b \otimes \chi \circ \theta^a) \circ \Delta(w)(v) + \\ &+ (\text{id} \otimes \chi \circ \theta^a \circ C^b) \circ \Delta(w)(v) + (\text{id} \otimes \chi \circ \theta^a \otimes \chi \circ \theta^b) \circ (\Delta \otimes \text{id}) \circ \Delta(w)(v). \end{aligned}$$

Since χ is a representation we have $(\chi \circ \theta^a \otimes \chi \circ \theta^b - \pm \chi \circ \theta^b \otimes \chi \circ \theta^a) \circ \Delta = \chi \circ [\theta^a, \theta^b]$. Since Δ is associative and C a representation we have

$$\begin{aligned} & [\Theta^a, \Theta^b](w \otimes v) = \\ &= C^{[a,b]}(w) \otimes v + (\text{id} \otimes \chi \circ (\theta^a \circ C^b - \pm \theta^b \circ C^a)) \circ \Delta(w)(v) + \\ &+ (\text{id} \otimes \chi \circ [\theta^a, \theta^b]) \circ \Delta(w)(v). \end{aligned}$$

This formula shows that Θ is a representation if and only if

$$(127) \quad (\text{id} \otimes \chi \circ (\theta^a \circ C^b - \pm \theta^b \circ C^a + [\theta^a, \theta^b] - \theta^{[a,b]})) \circ \Delta = 0 \quad \forall a, b \in \mathfrak{g}.$$

To show that the direct part assertion of the lemma is a necessary condition, it is enough to compose $\delta \otimes \text{id}_{U(\mathfrak{g})}$ with the left hand side of (127).

ii) To show the direct part is sufficient to consider a faithful representation χ of $\mathfrak{h}$, for instance the left regular representation in $U(\mathfrak{h})$. $\square$

Theorem 4.4. *The map $\Theta_{c,\chi,h,q,\mathfrak{g},\sigma}$ is a representation for any symmetric pair $(\mathfrak{g}, \sigma)$ and any representation χ of $\mathfrak{h}$, if and only if the functional equations*

$$(128) \quad -h(t+u) + h(t) + h(u) - h(t)h(u) = 0,$$

$$(129) \quad \frac{q(t+u) - q(t)}{u} p_c(u) + \frac{q(t+u) - q(u)}{t} p_c(t) - q(t)q(u) + h(t+u) = 0,$$

$$(130) \quad q(t) + \frac{h(t+u) - h(u)}{t} p_c(t) - q(t)h(u) = 0,$$

are satisfied in $\mathbb{K}_0[[t, u]]$.

Proof. Let a and b in $\mathfrak{g}$. For $X \in \mathfrak{go}$, $i, j \in \mathbb{N}$, we set

$$(t^i u^j : [a, b])_X = [(\text{ad } X)^i a, (\text{ad } X)^j b].$$

We use the same notation for the Lie algebra $F(\mathfrak{q}, \mathfrak{g})$. Thus, for a and b in $\mathfrak{g}$, and the generic point $y \in F(\mathfrak{q}, \mathfrak{g})$ we get an element $(t^i u^j : [a, b])_y \in F(\mathfrak{q}, \mathfrak{g})$. Let $k(t, u) = \sum k_{ij} t^i u^j \in \mathbb{K}_0[[t, u]]$. We define

$$(k(t, u) : [a, b])_y := \sum k_{ij} (t^i u^j : [a, b])_y \in F(\mathfrak{q}, \mathfrak{g}).$$

This is possible because this sum is locally finite.

Let us denote $h^b(t) = h(t)$ if $b \in \mathfrak{h}$, $h^b(t) = q(t)$ if $b \in \mathfrak{q}$, $p_c^b(t) = p_c(t)$ if $b \in \mathfrak{q}$, and $p_c^b(t) = -t$ if $b \in \mathfrak{h}$. We have (see [18])

$$\begin{aligned} \theta^a \circ C_c^b &= \left(-\frac{h^a(t+u) - h^a(t)}{u} p_c^b(u) : [a, b] \right)_y = \left(\frac{h^a(t+u) - h^a(u)}{t} p_c^b(t) : [b, a] \right)_y \\ [\theta^a, \theta^b] &= (h^a(t)h^b(u) : [a, b])_y \\ \theta^{[a, b]} &= (h^{[a, b]}(t+u) : [a, b])_y. \end{aligned}$$

Lemma 4.2 shows that Θ_χ is a representation of $\mathfrak{g}$ for all representations χ of $\mathfrak{h}$ if and only if

$$\begin{aligned} (h(t) - h(t+u) + h(u) - h(t)h(u) : [a, b])_y &= 0 \quad \forall a, b \in \mathfrak{h}, \\ \left(\frac{q(t+u) - q(t)}{u} p_c(u) + \frac{q(t+u) - q(u)}{t} p_c(t) + \right. \\ &\quad \left. - q(t)q(u) + h(t+u) : [a, b] \right)_y = 0 \quad \forall a, b \in \mathfrak{q}, \\ \left(\frac{h(t+u) - h(t)}{u} p_c(u) + q(u) - h(t)q(u) : [a, b] \right)_y &= 0 \quad \forall a \in \mathfrak{h}, b \in \mathfrak{q}, \\ \left(q(t) + \frac{h(t+u) - h(u)}{t} p_c(t) - q(t)h(u) : [a, b] \right)_y &= 0 \quad \forall a \in \mathfrak{q}, b \in \mathfrak{h}. \end{aligned}$$

The last two equations are identical. As in [18] we see that these equations are verified for any Lie symmetric pair $(\mathfrak{g}, \sigma)$ if and only if the functional equations in the statement of the theorem are satisfied. $\square$

We study the functional equations of Theorem 4.4.

Lemma 4.3. *i) Equation (128) has an unique solution $h(t) \in \mathbb{K}_0[[t^2]]$ such that $h(0) = 0$. It is $h = 0$.*

ii) Equation (128) has an unique solution $h(t) \in \mathbb{K}_0[[t^2]]$ such that $h(0)$ is invertible. It is $h = 1$.

Proof. *i)* If $h(0) = 0$ we can divide the equation (128) by t getting $\frac{-h(t+u)+h(u)}{t} = \frac{h(t)}{t}(h(u)-1)$. The evaluation at $t = 0$ gives $-h'(u) = h'(0)(h(u)-1)$. As $h(u) = h(-u)$ we have $h'(0) = 0$, so $h'(u) = 0$ and $h(u)$ is constant. This proves *i*.

ii) Evaluating at $t = 0$ we get $0 = h(0)(h(u)-1)$. If $h(0)$ is invertible, the unique solution is $h(u) = 1$. $\square$

Remark 4.5. *If $h = 0$, equation (130) gives $q = 0$. In this case, Θ_χ does not depend on χ . It is the tensor product of the representation C_c in $S(\mathfrak{q})$ with the trivial representation of $\mathfrak{g}$ in the module V .*

We now consider the more interesting case $h = 1$. Equations (128) and (130) are satisfied, and equation (129) becomes

$$(131) \quad \frac{q(t+u) - q(t)}{u} p_c(u) + \frac{q(t+u) - q(u)}{t} p_c(t) = q(t)q(u) - 1.$$

We introduce

$$(132) \quad q_c(t) = -\operatorname{th}\left(\frac{t}{2c}\right) = -2 \sum_{n=1}^{\infty} b_{2n} \frac{(2^{2n} - 1)t^{2n-1}}{c^{2n-1}(2n)!} = -\frac{1}{2c}t + \frac{1}{24c^3}t^3 + \dots$$

Remark 4.6. The Taylor series of $\operatorname{th}(t)$ is of course well known. However, since it plays a role in the proof of theorem 4.6 below, we remark that (132) follows from (114) and the relation

$$(133) \quad q_c(2t) = \frac{p_c(t) - p_c(2t)}{t},$$

which is equivalent to the identity $\operatorname{th}(t) + \coth(t) - 2\coth(2t) = 0$.

Proposition 4.1. The function q_c is the unique solution $q \in t\mathbb{K}_0[[t]]$ of the functional equation (131).

Proof. Since $q(0) = 0$, the evaluation at $u = 0$ of (131) gives

$$q'(t)c + \frac{q(t)}{t}p_c(t) \equiv q'(t)c + q(t)\coth\left(\frac{t}{c}\right) = -1.$$

This differential equation has exactly one solution such that $q(0) = 0$, because the equation recursively determines all the coefficients of q . For instance, for the first one we obtain $q_1c + q_1c = -1$ and $q_1 = -\frac{1}{2c}$, which is compatible with (132). This solution is of course q_c , and one checks that q_c also satisfy the functional equation (131). $\square$

We summarize the main result of this section. Let $c \in \mathbb{K}_0^\times$. We denote by $\mathfrak{g} \ni a \rightarrow \theta_c^a \in F(\mathfrak{q}, \mathfrak{h})$ the linear map such that

$$(134) \quad \theta_c^a = a \text{ for } a \in \mathfrak{h}, \text{ and } \theta_c^a = q_c(\operatorname{ad} y)(a) \text{ for } a \in \mathfrak{q}.$$

We use the notation $\Theta_{c,\chi} = \Theta_{c,\chi,1,q_c,\mathfrak{g},\sigma}$ (see formula (125)). For $a \in \mathfrak{h}$, $\Theta_{c,\chi}^a$ does not depend on c . It is the natural endomorphism of $S(\mathfrak{q}) \otimes V$ coming from the adjoint action of a in $S(\mathfrak{q})$ and from the action $\chi(a)$ in V . For $a \in \mathfrak{q}$, we have

$$(135) \quad \Theta_{c,\chi}^a = C_c^a \otimes \operatorname{id}_V + (\operatorname{id}_{S(\mathfrak{q})} \otimes \chi \circ \theta_c^a) \circ \Delta_{S(\mathfrak{q})}.$$

Corollary 4.1. i) $\Theta_{c,\chi}$ is a representation of $\mathfrak{g}$ in $S(\mathfrak{q}) \otimes V$.

ii) $I_c \otimes \operatorname{id}_V$ intertwines $\Theta_{1,\chi}$ and $\Theta_{c,\chi}$.

We use the notations of the beginning of this section. The following theorem contains theorem 4.3 as a particular case. We recall the notations $\alpha_a, \alpha_1^a, \theta_a, \theta_1^a$ from proposition 3.1 and formulas (116), (134).

Theorem 4.5. i) For $a \in \mathfrak{g}$, we have $\bar{l}_{\chi,a} = \Theta_{1,\chi}^a$.

ii) For $a \in \mathfrak{g}$, we have $\alpha_a = \alpha_1^a$ and $\theta_a = \theta_1^a$.

Proof. i) We extend $\Theta_{1,\chi}$ to a map from $U(\mathfrak{g})$ to $\operatorname{End}(S(\mathfrak{q}) \otimes V)$. Since $h = 1$ it induces the well defined map $\operatorname{map} \tau : U(\mathfrak{g}) \otimes_{\mathfrak{h}} V \ni u \otimes v \rightarrow \Theta_{1,\chi}^u(1 \otimes v) \in S(\mathfrak{q}) \otimes V$ (recall formula (126)). By definitions, τ intertwines the left multiplication in $U(\mathfrak{g}) \otimes_{\mathfrak{h}} V$ and the representation $\Theta_{1,\chi}$ in $S(\mathfrak{q}) \otimes V$. As in the proof of theorem 4.3, we see that $\tau \circ (\beta \otimes \operatorname{id}_V)$ is the identity map of $S(\mathfrak{q}) \otimes V$.

ii) It is a rephrasing of i). $\square$

Remark 4.7. When $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is a Lie superalgebra over a field of characteristic zero, and $\mathfrak{h} = \mathfrak{g}_0$, the formula $\bar{l}_{\chi,a} = \Theta_{1,\chi}^a$ is due to Koszul [13]. For real analytic symmetric space, the function $\operatorname{th}(\frac{t}{2})$ occurs in [22].

We get a similar result for the coinduced representation $F(G/H, V)$, and the corresponding representation in $F(\mathfrak{q}, V)$. Indeed: from (80) and theorem 4.5 we obtain the following corollary.

Corollary 4.2. *For $a \in \mathfrak{g}$, we have*

$$(136) \quad -\bar{\zeta}_{\chi, a} = -\zeta_{\alpha_1^a} + \chi(\theta_1^a).$$

4.3. The Jacobian of the exponential map for a symmetric pair. In this subsection, we assume that $\mathfrak{q}$ is free of finite rank and we compute the Jacobian of the exponential map $\mathfrak{q} \rightarrow G/H$, as defined by formula (97).

Let $(e_i)_{1 \leq i \leq n}$ be a basis of $\mathfrak{q}$, and $(x^i)_{1 \leq i \leq n}$ be the dual basis of $\mathfrak{q}^*$. Then $F(\mathfrak{q}) = \mathbb{K}[[x^1, \dots, x^n]]$. The generic point y of $\mathfrak{q}$ is equal to $y = \sum e_i x^i \in F(\mathfrak{q}, \mathfrak{q}) = \mathfrak{q} \otimes F(\mathfrak{q})$. Since $F(\mathfrak{q}, \mathfrak{g}) = F(\mathfrak{q}, \mathfrak{h}) \oplus F(\mathfrak{q}, \mathfrak{q})$, we consider y as an element of the Lie $F(\mathfrak{q})$ -superalgebra $F(\mathfrak{q}, \mathfrak{g})$.

If $k \in 2\mathbb{N}$, $\text{ad}^k(y)$ stabilizes $\mathfrak{q} \otimes F(\mathfrak{q})$ and we denote by $\text{str}_{\mathfrak{q}}(\text{ad}^k(y)) \in F(\mathfrak{q})_{\mathbf{0}}$ its supertrace, which is an homogeneous polynomial of degree k in the x^i . More generally, for $w(t) = w_0 + w_2 t^2 + \dots \in \mathbb{K}_{\mathbf{0}}[[t^2]]$, $\text{str}_{\mathfrak{q}}(w(\text{ad } y))$ is defined and we have the Taylor expansion

$$(137) \quad \text{str}_{\mathfrak{q}}(w(\text{ad } y)) = \sum_{k \geq 0} w_k \text{str}_{\mathfrak{q}}(\text{ad}^k(y)) \in F(\mathfrak{q})_{\mathbf{0}}.$$

Similarly, for $r(t) = 1 + r_2 t^2 + \dots \in \mathbb{K}_{\mathbf{0}}[[t^2]]$, $r(\text{ad } y)$ induces an even invertible $F(\mathfrak{q})$ -linear transformation of $\mathfrak{q} \otimes F(\mathfrak{q})$. We denote by $\text{Ber}_{\mathfrak{q}}(r(\text{ad } y))$ its Berezinian.

For $c \in \mathbb{K}_{\mathbf{0}}^*$ we define

$$(138) \quad J_c := \text{Ber}_{\mathfrak{q}} \left(\frac{\text{sh}(\text{ad} \frac{y}{c})}{\text{ad} \frac{y}{c}} \right) \in F(\mathfrak{q})_{\mathbf{0}},$$

and

$$(139) \quad w_c(t) = \log \left(\frac{\text{sh}(\frac{t}{c})}{\frac{t}{c}} \right) \in t^2 \mathbb{K}_{\mathbf{0}}[[t^2]].$$

Because of formula (114) and

$$(140) \quad w'_c(t) = \frac{1}{ct} \left(t \coth \left(\frac{t}{c} \right) - c \right) = \frac{1}{p_c(0)} \frac{p_c(t) - p_c(0)}{t},$$

we have

$$(141) \quad w_c(t) = \sum_{n=1}^{\infty} b_{2n} \frac{2^{2n} t^{2n}}{c^{2n} 2n(2n)!} = \frac{1}{6c^2} t^2 - \frac{1}{180c^4} t^4 + \dots$$

Because of (30) we obtain

$$(142) \quad J_c = \exp(\text{str}_{\mathfrak{q}}(w_c(\text{ad } y))).$$

We write the beginning of the resulting Taylor expansion of J_c :

$$(143) \quad J_c = 1 + \frac{1}{6c^2} \text{str}_{\mathfrak{q}}(\text{ad}^2(y)) - \frac{1}{180c^4} \text{str}_{\mathfrak{q}}(\text{ad}^4(y)) + \frac{1}{72c^4} \text{str}_{\mathfrak{q}}^2(\text{ad}^2(y)) + \dots$$

We start with some computations on supertraces needed in the proof of theorem 4.6. For $a \in \mathfrak{q}$ and $i, j \in \mathbb{N}$ such that $i+j \in 2\mathbb{N}+1$, the element $\text{ad}^i(y) \text{ad } a \text{ad}^j(y)$ of $\text{End}_{F(\mathfrak{q})}(\mathfrak{g} \otimes F(\mathfrak{q}))$ stabilizes $\mathfrak{q} \otimes F(\mathfrak{q})$. Let t and u be even formal variables. We use the notation

$$(144) \quad S(t^i u^j : a) = \text{str}_{\mathfrak{q}}(\text{ad}^i(y) \text{ad } a \text{ad}^j(y)).$$

Let $k(t, u) = \sum_{i, j \geq 0} k_{ij} t^i u^j \in \mathbb{K}[[t, u]]$ be such that $k_{ij} = 0$ if $i+j \in 2\mathbb{N}$. We define the locally finite sum

$$(145) \quad S(k(t, u) : a) = \sum_{i, j \geq 0} k_{ij} S(t^i u^j : a) \in F(\mathfrak{q}).$$

For example, for $q(t) \in t\mathbb{K}_0[[t^2]]$ we have $q(\text{ad } y)(a) \in F(\mathfrak{q}, \mathfrak{h})$, and $\text{ad}(q(\text{ad } y)(a))$ stabilizes $\mathfrak{q} \otimes F(\mathfrak{q})$. We have

$$(146) \quad \text{str}_{\mathfrak{q}}(\text{ad}(q(\text{ad } y)(a))) = S(q(t-u) : a).$$

Lemma 4.4. $S((t^2 - u^2)k(t, u) : a) = 0$.

Proof. It is sufficient to prove it for a monomial $k = t^i u^j$. Let Y and X be the restrictions of $\text{ad}^2(y)$ and $\text{ad}^i(y) \text{ad } a \text{ad}^j(y)$ to $\mathfrak{q} \otimes F(\mathfrak{q})$. Since Y is even, we have $\text{str}_{\mathfrak{q}}(YX) = \text{str}_{\mathfrak{q}}(XY)$ which proves the lemma. $\square$

Let $g(t) = g_0 + g_2 t^2 + \dots \in \mathbb{K}_0[[t^2]]$. We consider the vector field $g(\text{ad } y)(a)$ on $\mathfrak{q}$.

Proposition 4.2. *We have*

$$\text{div}(\zeta_{g(\text{ad } y)(a)}) = S\left(-\frac{g(t) - g(t-u)}{u} : a\right).$$

Proof. It is sufficient to prove the statement for a monomial $g(t) = t^n$ with n even. We argue like in the proof of proposition 2.2, replacing x with y , and str with $\text{str}_{\mathfrak{q}}$. Proposition 4.2 is equivalent to formula (54). $\square$

Let $r(t) = r_0 + r_2 t^2 + \dots \in \mathbb{K}_0[[t^2]]$. We consider the function $\text{str}_{\mathfrak{q}}(r(\text{ad } y)) \in F(\mathfrak{q})$.

Proposition 4.3. *We have*

$$\zeta_{g(\text{ad } y)(a)}(\text{str}_{\mathfrak{q}}(r(\text{ad } y))) = S\left(\frac{r(t) - r(u)}{t-u} g(t-u) : a\right).$$

Proof. It is sufficient to prove the statement for a monomial $r(t) = t^n$ with n even. We argue like in the proof of proposition 2.3, replacing x with y , and str with $\text{str}_{\mathfrak{q}}$. Proposition 4.2 is equivalent to formula (59). $\square$

Lemma 4.5. *Let $c \in \mathbb{K}_0^\times$. For all $a \in \mathfrak{g}$ we have*

$$(147) \quad \zeta_{\alpha_c^a}(\text{str}_{\mathfrak{q}}(w_c(\text{ad } y))) + \text{div } \zeta_{\alpha_c^a} - \text{str}_{\mathfrak{q}}(\theta_c^a) = 0.$$

Proof. We first consider the case $a \in \mathfrak{q}$. From the definition (116) of α_c^a and propositions 4.3 and 4.2, we obtain

$$(148) \quad \zeta_{\alpha_c^a}(\text{str}_{\mathfrak{q}}(w_c(\text{ad } y))) = S\left(\frac{w_c(t) - w_c(u)}{t-u} p_c(t-u) : a\right).$$

and

$$(149) \quad \text{div } \zeta_{\alpha_c^a} = S\left(-\frac{p_c(t) - p_c(t-u)}{u} : a\right).$$

From the definition (134) of θ_c^a and equation (146) we obtain

$$(150) \quad -\text{str}_{\mathfrak{q}}(\theta_c^a) = -S(q_c(t-u) : a).$$

Thus, the left hand side of (147) is equal to $S(s(t, u) : a)$ with

$$s(t, u) := \frac{w_c(t) - w_c(u)}{t-u} p_c(t-u) - \frac{p_c(t) - p_c(t-u)}{u} - q_c(t-u).$$

Lemma 4.4 shows that to prove (147), it is sufficient to prove that $s(t, u)$ belongs to the ideal $(t^2 - u^2)\mathbb{K}_0[[t, u]]$. For this, it is sufficient to prove that we have $s(t, t) = 0$ and $s(t, -t) = 0$. We have (see (141))

$$(151) \quad s(t, t) = w'_c(t)p_c(0) - \frac{p_c(t) - p_c(0)}{t} \text{ and } s(t, -t) = \frac{p_c(t) - p_c(2t)}{t} - q_c(2t).$$

Strangely enough, we already noticed the relation $s(t, t) = 0$ in formula (140) and the relation $s(t, -t) = 0$ in (133).

Last, we consider the simpler case $a \in \mathfrak{h}$. The proof is the same with $p_c(t)$ replaced by $-t$, and $q_c(t)$ replaced by 1. We have $s(t, t) = -\frac{-t}{t} - 1 = 0$ and $s(t, -t) = \frac{-t+2t}{t} - 1 = 0$. $\square$

Theorem 4.6. *Let $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$ be a symmetric pair such that $\mathfrak{q}$ is free of finite rank. The Jacobian $J \in F(\mathfrak{q})$ of the exponential map $\mathfrak{q} \rightarrow G/H$ is given by the formula $J = J_1$.*

Proof. We verify that J_1 satisfies the conditions of lemma 3.3. We already showed that $\delta(J_1) = 1$ in formula (143). Let $c \in \mathbb{K}_{\mathbf{0}}^*$. For a vector field $\alpha \in F(\mathfrak{q}, \mathfrak{q})$ we get from (30) the relation

$$\frac{\zeta_\alpha(J_c)}{J_c} = \zeta_\alpha(\text{str}_{\mathfrak{q}}(w_c(\text{ad } y))).$$

Let $a \in \mathfrak{g}$. We recall the notations $\alpha_c^a \in F(\mathfrak{q}, \mathfrak{q})$ and $\theta_c^a \in F(\mathfrak{q}, \mathfrak{h})$ in theorem 4.5. The second condition (98) of lemma 3.3 follows from lemma 4.5 choosing $c = 1$. $\square$

4.4. Gorelik elements. We recall the definition (1) of the twisted adjoint representation ad' of $\mathfrak{g}$ in $U(\mathfrak{g})$. We extend ad' to a representation of $U(\mathfrak{g})$ in $U(\mathfrak{g})$ and we consider the map $\tilde{\gamma} : U(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ defined by

$$(152) \quad \tilde{\gamma}(u) = \text{ad}'(u)(1) \text{ for } u \in U(\mathfrak{g}).$$

Remark 4.8. *By definition the map $\tilde{\gamma}$ intertwines the left regular representation and the twisted adjoint action.*

Moreover, since $\text{ad}'(a)(1) = \text{ad}(a)(1) = 0$ for $a \in \mathfrak{h}$, the map $\tilde{\gamma}$ gives a map

$$(153) \quad \gamma : U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h} \rightarrow U(\mathfrak{g}).$$

Lemma 4.6. (Gorelik [9]). *We have $\gamma \circ \beta|_{S(\mathfrak{q})} = \beta \circ I_2$.*

For example, for $b \in \mathfrak{q}$, we have $\gamma(b) = \text{ad}'(b)(1) = b1 - 1\sigma(b) = b + b = 2b$. Thus γ is an isomorphism of coalgebras from $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$ to $\beta(S(\mathfrak{q}))$.

Proposition 4.4. i) *The submodule $\beta(S(\mathfrak{q})) \subset U(\mathfrak{g})$ is stable under the twisted adjoint action;*
 ii) *as a $\mathfrak{g}$ -module it is isomorphic to $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$;*
 iii) *the map $\beta : S(\mathfrak{q}) \rightarrow \beta(S(\mathfrak{q}))$ intertwines the representation C_2 with the twisted adjoint action.*

Remark 4.9. *The first assertion is due to [1], the second to [9], and the last one follows from remark 4.8, lemma 4.6, theorem 4.3, and remark 4.2.*

We assume moreover that the dimension of $\mathfrak{q}$ is $(0, q)$ and that the unimodularity condition (10) holds. Let $(e_i)_{1 \leq i \leq q}$ be a totally ordered basis of $\mathfrak{q}$, and (x^i) the dual basis of $\mathfrak{q}^*$. We have $F(\mathfrak{q}) = \mathbb{K}[x^1, \dots, x^q]$ and the formal sum (143) which defines J_c is finite. We recall that, by theorems 3.1 and 4.6, the element

$$\tilde{\beta}(J_1 e_1 \dots e_q) \in U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$$

is basis of the $\mathbb{K}$ -module of $\mathfrak{g}$ -invariant elements of $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$, and a basis of $U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}$ as a $F(G/H)$ -module.

Theorem 4.7. *Let $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$ be a symmetric pair such that $\mathfrak{q}$ is free of finite rank $(0, q)$ and such that the unimodularity condition (10) holds. Let $(e_i)_{1 \leq i \leq q}$ be a totally ordered basis of $\mathfrak{q}$. The element $\beta(J_2 e_1 \dots e_q) \in U(\mathfrak{g})$ is a basis of the $\mathbb{K}$ -module of the invariants of the twisted adjoint action of $\mathfrak{g}$ in $\beta(S(\mathfrak{q}))$.*

Proof. Recall that $J_1 e_1 \dots e_q$ is an invariant for the representation $\mathfrak{g} \ni a \rightarrow C_1^a$ (see theorem 4.3, ii). Then the theorem follows from proposition 4.4 part iii, remark 4.2, and $I_c(J_1 e_1 \dots e_q) = c^q J_c e_1 \dots e_q$. $\square$

The following corollary is the particular case of theorem 4.7 corresponding to $\mathbb{K} = \mathbb{K}_0$, $\mathfrak{h} = \mathfrak{g}_0$, and $\mathfrak{q} = \mathfrak{g}_1$ free of finite rank. We state it in the conditions of the introduction.

Corollary 4.3. *Let $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ be a Lie superalgebra of finite dimension (p, q) over a field $\mathbb{K}$ of characteristic 0, and $d \in S^q(\mathfrak{g}_1)$ with $d \neq 0$. Assume that the unimodularity assumption (2) holds. Then $\beta(J_2 d)$ is a Gorelik element of $U(\mathfrak{g})$.*

REFERENCES

- [1] D. Arnaudon, M. Bauer, and L. Frappat, *On Casimir's ghost*, Comm. Math. Phys. **187** (1997), no. 2, 429–439.
- [2] Allen D. Bell and Rolf Farnsteiner, *On the theory of Frobenius extensions and its application to Lie superalgebras*, Trans. Amer. Math. Soc. **335** (1993), no. 1, 407–424.
- [3] F. A. Berezin and G. I. Kac, *Lie groups with commuting and anticommuting parameters*, Mat. Sb. (N.S.) **82** (124) (1970), 343–359 (Russian).
- [4] Robert J. Blattner, *Induced and produced representations of Lie algebras*, Trans. Amer. Math. Soc. **144** (1969), 457–474.
- [5] Sophie Chemla, *Cohomologie locale de Grothendieck et représentations induites de superalgèbres de Lie*, Math. Ann. **297** (1993), no. 2, 371–382 (French).
- [6] ———, *Propriétés de dualité dans les représentations coinduites de superalgèbres de Lie*, Ann. Inst. Fourier (Grenoble) **44** (1994), no. 4, 1067–1090 (French, with English and French summaries).
- [7] ———, *Poincaré duality for k -A Lie superalgebras*, Bull. Soc. Math. France **122** (1994), no. 3, 371–397.
- [8] P. M. Cohn, *A remark on the Birkhoff-Witt theorem*, J. London Math. Soc. **38** (1963), 197–203.
- [9] Maria Gorelik, *On the ghost centre of Lie superalgebras*, Ann. Inst. Fourier (Grenoble) **50** (2000), no. 6, 1745–1764 (2001).
- [10] ———, *Strongly typical representations of the basic classical Lie superalgebras*, J. Amer. Math. Soc. **15** (2002), no. 1, 167–184 (electronic).
- [11] Maria Gorelik and Emmanuel Lanzmann, *The annihilation theorem for the completely reducible Lie superalgebras*, Invent. Math. **137** (1999), no. 3, 651–680.
- [12] Yvette Kosmann-Schwarzbach and Juan Monterde, *Divergence operators and odd Poisson brackets*, Ann. Inst. Fourier **52** (2002), no. 2, 419–456.
- [13] J.-L. Koszul, *Graded manifolds and graded Lie algebras*, (Florence, 1982), Pitagora, Bologna, 1983, pp. 71–84.
- [14] Andrzej Lesniewski, *A remark on the Casimir elements of Lie superalgebras and quantized Lie superalgebras*, J. Math. Phys. **36** (1995), no. 3, 1457–1461.
- [15] Yuri I. Manin, *Gauge field theory and complex geometry*, 2nd ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 289, Springer-Verlag, Berlin, 1997.
- [16] Ian M. Musson, *On the center of the enveloping algebra of a classical simple Lie superalgebra*, J. Algebra **193** (1997), no. 1, 75–101.
- [17] Emanuela Petracchi, *Functional equations and Lie algebras*, Tesi di Dottorato, Università di Roma “La Sapienza”, 2003 (English).
- [18] ———, *Universal representations of Lie algebras by coderivations*, Bull. Sci. Math. **127** (2003), no. 5, 439–465.
- [19] Georges Pinczon, *The enveloping algebra of the Lie superalgebra $\mathfrak{osp}(1, 2)$* , J. Algebra **132** (1990), no. 1, 219–242.
- [20] David E. Radford, *Divided power structures on Hopf algebras and embedding Lie algebras into special-derivation algebras*, J. Algebra **98** (1986), no. 1, 143–170.
- [21] François Rouvière, *Espaces symétriques et méthode de Kashiwara-Vergne*, Ann. Sci. École Norm. Sup. (4) **19** (1986), no. 4, 553–581 (French, with English summary).
- [22] ———, *Fibrés en droites sur un espace symétrique et analyse invariante*, J. Funct. Anal. **124** (1994), no. 2, 263–291 (French, with English and French summaries).
- [23] Manfred Scheunert, *The theory of Lie superalgebras*, Lecture Notes in Mathematics, vol. 716, Springer, Berlin, 1979.
- [24] T. Voronov, *Geometric integration theory on supermanifolds*, Soviet Scientific Reviews, Section C: Mathematical Physics Reviews, vol. 9, Harwood Academic Publishers, Chur, 1991.
- [25] Alexander A. Voronov, Yu. I. Manin, and I. B. Penkov, *Elements of supergeometry*, Current problems in mathematics. Newest results, Vol. 32, Itogi Nauki i Tekhniki, Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1988, pp. 3–25 (Russian). Translated in J. Soviet Math. **51** (1990), no. 1, 2069–2083.

UNIVERSITY DENIS DIDEROT-PARIS 7, INSTITUT DE MATHÉMATIQUES DE JUSSIEU, C.P. 7012, 2 PLACE
JUSSIEU, F-75251 PARIS CEDEX 05

E-mail address: `duflo@math.jussieu.fr`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, 40 ST. GEORGE ST. TORONTO, ONTARIO,
CANADA M5S 2E4

E-mail address: `petracci@math.toronto.edu`